



Penn Medicine

Perelman School of Medicine
University of Pennsylvania Health System

Penn Diabetes Research Center MOUSE PHENOTYPING, PHYSIOLOGY & METABOLISM CORE

Director: Joseph Baur, Ph.D.

Technical Director: Jennifer Rojas, Ph.D.

Request for Clinical Chemistry Analysis

Lab/PI:	Requestor:
Phone:	Request Date:
Email:	Account Number:

Select	Species
<input type="checkbox"/>	Mouse
<input type="checkbox"/>	Rat
<input type="checkbox"/>	Other (inquire)

IDEXX VetTEST Chemistry Analyzer

Sample volume: The VetTest Chemistry Analyzer requires a minimum of 40 µl of plasma or serum sample (refer to sample preparation below) to run one test plus 10 µl for each additional test.

Preparing a plasma sample: We recommend using a lithium-heparin tube with a gel barrier. Separate plasma from red cells within five minutes when using lithium-heparin.

1. Use the appropriate tube. Do not use EDTA.
2. Use the appropriate sample collection device.
3. Draw the sample gently. Please avoid hemolysis. Use the correct blood-to-lithium-heparin ratio. Gently invert the sample for 30 seconds to mix.
4. Centrifuge the samples at least 120 seconds at a minimum of 12,000 Relative Centrifugal Force (RCF).
5. Transfer the plasma sample into a fresh 1.5mL micro-centrifuge tube and store at -20°C for subsequent assay analysis (sample should be analyzed within 1 week).

Preparing a serum sample:

1. Use the appropriate tube.
2. Use the appropriate sample collection device.
3. Draw the sample gently. Please avoid hemolysis. Let the sample clot for a minimum of 20 minutes at room temperature.
4. Centrifuge the samples at least 120 seconds at a minimum of 12,000 Relative Centrifugal Force (RCF).
5. Transfer the plasma sample into a fresh 1.5mL micro-centrifuge tube and store at -20°C for subsequent assay analysis (sample should be analyzed within 1 week).
- 6.

Select	Diagnostic Panel Assays	No. Of Samples	Unit Price	Total Price
--------	-------------------------	----------------	------------	-------------

Liver/Gut Function:

<input type="checkbox"/>	Albumin Alkaline phosphatase Alanine aminotransferase Amylase Aspartate aminotransferase Gamma-glutamyltransferase Lactate dehydrogenase Lipase Total bilirubin Total protein		\$80/panel set/sample	
--------------------------	--	--	--------------------------	--

Kidney Function:



Penn Medicine

Perelman School of Medicine
University of Pennsylvania Health System

Penn Diabetes Research Center MOUSE PHENOTYPING, PHYSIOLOGY & METABOLISM CORE

Director: Joseph Baur, Ph.D.

Technical Director: Jennifer Rojas, Ph.D.

	Albumin Ammonia Calcium Creatinine Magnesium Inorganic phosphate Total protein Urea Uric acid		\$80/panel set/sample	
--	---	--	----------------------------------	--

Muscle Function:

	Albumin Calcium Creatine kinase Creatinine Lactate Lactate dehydrogenase Magnesium Inorganic phosphate Total protein Urea		\$80/panel set/sample	
--	--	--	----------------------------------	--

Total \$

Select	Individual Assays	No. Of Samples	Unit Price	Total Price
	Albumin		\$10/assay	
	Alkaline phosphatase		\$10/assay	
	Alanine aminotransferase		\$10/assay	
	Amylase		\$10/assay	
	Aspartate aminotransferase		\$10/assay	
	Ammonia		\$10/assay	
	Calcium		\$10/assay	
	Cholesterol		\$10/assay	
	Creatine kinase		\$10/assay	
	Creatinine		\$10/assay	
	Gamma-glutamyltransferase		\$10/assay	
	Glucose		\$10/assay	
	Lactate		\$10/assay	
	Lactate dehydrogenase		\$10/assay	
	Lipase		\$10/assay	
	Magnesium		\$10/assay	
	Inorganic phosphate		\$10/assay	
	Triglycerides		\$10/assay	
	Total bilirubin		\$10/assay	
	Total protein		\$10/assay	
	Urea		\$10/assay	
	Uric acid		\$10/assay	

Total \$



Penn Diabetes Research Center MOUSE PHENOTYPING, PHYSIOLOGY & METABOLISM CORE

Director: Joseph Baur, Ph.D.

Technical Director: Jennifer Rojas, Ph.D.

Quantitative Enzymatic-Colorimetric Assays

Specimen Collection and Preparation: Specimen may be serum, or plasma collected with EDTA as anticoagulant and stored at -20°C (short term) or -80°C (long term). Please avoid hemolysis. To avoid multiple freeze/thaw cycles, samples must be aliquoted into separate tubes for each assay.

Interfering substances for triglyceride assay: Blood collecting devices containing glycerol (glycerin) cannot be used. Gross hemolysis or high bilirubin values, will produce falsely elevated values.

Interfering substances for NEFA assay: Any specimen containing heparin is unsuitable for analysis.

Select	Individual Assays	Assay Manufacturer	Sample Type & Volume (for duplicate analysis)	No. Of Samples	Unit Price	Minimum Order	Total Price
	Total Cholesterol	Stanbio laboratory	Plasma or serum (10 µl)		\$5/assay	\$50	
	Triglycerides	Stanbio laboratory	Plasma or serum (10 µl)		\$5/assay	\$50	
	Non-esterified free fatty acids (inquire)	WAKO Diagnostics	Serum (10 µl)		\$5/assay	\$50	

Total \$

Sample Shipping and Delivery:

UPenn Investigators: Place samples in designated retainer bin in the drop-off freezer #2 in the 12-170 corridor.

External investigators: Please contact the Technical Director, Jennifer Rojas to arrange delivery of samples.

Sample Labeling:

The sample box and each tube should be uniquely labeled (e.g. Date, Sample ID, Investigator's name/initials, and type of assay requested).

Please send order request and all inquiries to the Technical Director, Jennifer Rojas at Jennifer.Rojas@penntmedicine.upenn.edu.

Kindly acknowledge the Penn Diabetes Research Center grant P30-DK19525, and the services of the Mouse Phenotyping, Physiology and Metabolism Core in all publications and presentations.