SHINE Shift Change Checksheet-Control

Please review time on the study protocol, clinical or safety events or other concerns as well as the items below. Page the Stroke Pager (215-452-2793) or call Melissa Kruszewski (215-264-4382) with questions or concerns about the study protocol.

Sh	ift change date/time/initials:,,	Notes:
	Have nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, study laptop and paper treatment log)?	
	Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)?	
	What level is the patient on in the sliding scale? Has the study team reviewed/advanced to higher level?	
	Is a new bag with study infusion due? a) Remember the IDS prepares this – bag expires: b) Is the infusion pump programmed with 'New guardrail drug' instead of insulin/saline?	
	Has there been neuroworsening that has persisted(NIHSS ≥4 points higher than prior NIHSS)? Has the study team been notified? a) Remember that NIHSS score is required 24 hrs from onset.	
	 Nutritional status a) Has there been a change from NPO to PO? Is a change anticipated? b) If PO or on bolus tube feeds, is 60 gram carbohydrate diet ordered? c) Remind nurse to hold lunch until 12:00 and hold dinner until 18:00 AFTER scheduled glucose check/SQ insulin dose. d) Is the patient on continuous tube feeds? IF TUBE FEEDS ARE PAUSED: STOP/HOLD study infusion as well 	
	Has the drip been paused or will it need to be for scheduled procedure? Has the study team been contacted?	
	If the patient might be discharged before 72 hours, has the study team been notified?	
	Have there been any problems with the study laptop or internet?	
Sh	ift change date/time/initials:,,	Notes:
Sh	ift change date/time/initials:,,,,	Notes:
_	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise,	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)? What level is the patient on in the sliding scale? Has the study team	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)? What level is the patient on in the sliding scale? Has the study team reviewed/advanced to higher level? Is a new bag with study infusion due? a) Remember the IDS prepares this – bag expires: b) Is the infusion pump programmed with 'New guardrail drug' instead of	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)? What level is the patient on in the sliding scale? Has the study team reviewed/advanced to higher level? Is a new bag with study infusion due? a) Remember the IDS prepares this – bag expires: b) Is the infusion pump programmed with 'New guardrail drug' instead of insulin/saline? Has there been neuroworsening that has persisted (NIHSS ≥4 points higher than prior NIHSS)? Has the study team been notified?	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)? What level is the patient on in the sliding scale? Has the study team reviewed/advanced to higher level? Is a new bag with study infusion due? a) Remember the IDS prepares this − bag expires: b) Is the infusion pump programmed with 'New guardrail drug' instead of insulin/saline? Has there been neuroworsening that has persisted (NIHSS ≥4 points higher than prior NIHSS)? Has the study team been notified? a) Remember that NIHSS score is required 24 hrs from onset. Nutritional status a) Has there been a change from NPO to PO? Is a change anticipated? b) If PO or on bolus tube feeds, is 60 gram carbohydrate diet ordered? c) Remind nurse to hold lunch until 12:00 and hold dinner until 18:00 AFTER scheduled glucose check/SQ insulin dose. d) Is the patient on continuous tube feeds? IF TUBE FEEDS ARE	Notes:
	Have the nurses been documenting finger sticks, drip rates, D50 administration and SQ insulin administration in all three places (Sunrise, the study laptop and on the paper treatment log)? Have there been episodes of hypoglycemia (<70mg/dL), severe hypoglycemia (<40mg/dL) or severe hyperglycemia (>500mg/dL)? What level is the patient on in the sliding scale? Has the study team reviewed/advanced to higher level? Is a new bag with study infusion due? a) Remember the IDS prepares this − bag expires: b) Is the infusion pump programmed with 'New guardrail drug' instead of insulin/saline? Has there been neuroworsening that has persisted (NIHSS ≥4 points higher than prior NIHSS)? Has the study team been notified? a) Remember that NIHSS score is required 24 hrs from onset. Nutritional status a) Has there been a change from NPO to PO? Is a change anticipated? b) If PO or on bolus tube feeds, is 60 gram carbohydrate diet ordered? c) Remind nurse to hold lunch until 12:00 and hold dinner until 18:00 AFTER scheduled glucose check/SQ insulin dose. d) Is the patient on continuous tube feeds? IF TUBE FEEDS ARE PAUSED: STOP/HOLD study infusion as well Has the drip been paused or will it need to be for scheduled procedure?	Notes: