

Featured Articles

Introduction to the recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease

Clifford R. Jack Jr.^{a,*}, Marilyn S. Albert^b, David S. Knopman^a, Guy M. McKhann^b,
Reisa A. Sperling^c, Maria C. Carrillo^d, Bill Thies^d, Creighton H. Phelps^e

^aDepartment of Radiology, Mayo Clinic, Rochester, MN, USA

^bDepartment of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

^cCenter for Alzheimer Research and Treatment, Brigham and Women's Hospital, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

^dAlzheimer's Association, Chicago, IL, USA

^eDivision of Neuroscience, National Institute on Aging, Bethesda, MD, USA

Abstract

Background: Criteria for the clinical diagnosis of Alzheimer's disease (AD) were established in 1984. A broad consensus now exists that these criteria should be revised to incorporate state-of-the-art scientific knowledge.

Methods: The National Institute on Aging (NIA) and the Alzheimer's Association sponsored a series of advisory round table meetings in 2009 whose purpose was to establish a process for revising diagnostic and research criteria for AD. The recommendation from these advisory meetings was that three separate work groups should be formed with each assigned the task of formulating diagnostic criteria for one phase of the disease: the dementia phase; the symptomatic, pre-dementia phase; and the asymptomatic, preclinical phase of AD.

Results: Two notable differences from the AD criteria published in 1984 are incorporation of biomarkers of the underlying disease state and formalization of different stages of disease in the diagnostic criteria. There was a broad consensus within all three workgroups that much additional work is needed to validate the application of biomarkers for diagnostic purposes. In the revised NIA-Alzheimer's Association criteria, a semantic and conceptual distinction is made between AD pathophysiological processes and clinically observable syndromes that result, whereas this distinction was blurred in the 1984 criteria.

Conclusions: The new criteria for AD are presented in three documents. The core clinical criteria of the recommendations regarding AD dementia and MCI due to AD are intended to guide diagnosis in the clinical setting. However, the recommendations of the preclinical AD workgroup are intended purely for research purposes.

© 2011 The Alzheimer's Association. All rights reserved.

Keywords:

Alzheimer's disease; Diagnostic criteria; Biomarkers; Imaging; Cerebrospinal fluid

1. Introduction

Criteria for the clinical diagnosis of Alzheimer's disease (AD) were established by the National Institute of Neurological and Communicative Disorders and Stroke (NINCDS) and the Alzheimer's Disease and Related Disorders Association (ADRDA) workgroup in 1984 [1]. These criteria were

universally adopted, have been extremely useful, and have survived intact without modification for more than a quarter of a century. However, in the intervening 27 years, important advances in our understanding of AD, in our ability to detect the pathophysiological process of AD, and changes in conceptualization regarding the clinical spectrum of the disease have occurred.

By 2009, broad consensus existed throughout academia and industry that the criteria should be revised to incorporate

*Corresponding author. Tel.: +1-507-284-9778; Fax: 1-507-284-2511.
E-mail address: jack.clifford@mayo.edu

scientific advances in the field. In response to this imperative, the National Institute on Aging (NIA) and the Alzheimer's Association sponsored a series of advisory round table meetings in 2009 whose purpose was to establish a process for revising diagnostic and research criteria for the continuum of AD. These advisory meetings included members from academia and industry with an international representation. The consensus from the advisory meetings was that three separate workgroups should be formed under the auspices of the NIA and Alzheimer's Association. One workgroup was assigned the task of formulating diagnostic criteria for the dementia phase of AD. A second was asked to focus on diagnostic criteria for the symptomatic pre-dementia phase of AD. The third was asked to propose a research agenda for the asymptomatic, preclinical phase of AD. Individuals were selected to serve in these workgroups, by the NIA and the Alzheimer's Association, with the objective of having balanced expertise and international representation from academia and industry. Between early and mid 2010, the three workgroups met through conference calls and in person (as feasible). Each formulated a set of recommendations. These recommendations were presented in a symposium at the 2010 International Conference on Alzheimer's Disease meeting. They were posted on the Alzheimer's Association Web site for a period of public comment over the summer of 2010. Comments received during this period, from the Web site and other venues, were given to the individual workgroups and incorporated into revisions of each document in the fall of 2010. Finally, a subcommittee representing individuals from each of the workgroups was asked to review the recommendations, particularly with regard to the approach to biomarkers, and a final round of revisions was made to each document in the late fall of 2010 for purposes of harmonizing the discussion of biomarkers. The final documents were submitted simultaneously in early 2011 to the NIA for review and to the *Journal* for peer review.

The charge to the workgroups was very specific and did not include several related topics. First, it was decided at the outset that a fourth, separate workgroup would be organized to develop revised pathological criteria. Thus, although neuropathologists were represented by each of the three workgroups, the recommendations of the three workgroups do not include a detailed discussion of neuropathologic criteria. The deliberations of the neuropathology workgroup are expected to appear later in 2011. Second, the workgroups were asked to outline future issues that need to be addressed by the research community as a whole, but the specifics for how this will be done, including potential timelines, are not included in these recommendations. This is particularly relevant to the discussion of biomarkers in each of the three documents. There was a consensus among the members of the workgroups that it was premature to define specific cut-points denoting normal versus abnormal values for the biomarkers discussed, and that much work remains to be done with regard to uniform assessment

and standardization of biomarkers. Third, the workgroups were specifically asked to focus on the spectrum of AD, and not to try to revise criteria for other neurodegenerative diseases or cerebrovascular dementias. Thus, the set of recommendations presented in this article only refer to other disorders because they relate to differential diagnosis of AD.

2. Historical background

The original NINCDS–ADRDA criteria rest on the notion that AD is a *clinical-pathological* entity [1]. The criteria were designed with the expectation that in most cases, subjects who met the clinical criteria would have AD pathology as the underlying etiology if the subject were to be presented for an autopsy. When the NINCDS–ADRDA criteria were formulated, it was believed that AD, like many other brain diseases, always exhibited a close correspondence between clinical symptoms and the underlying pathology, such that (1) AD pathology and clinical symptoms were synonymous, and (2) individuals either had fully developed AD pathology, in which case they were demented, or they were free of AD pathology, in which case they were not demented (at least not because of AD). However, in the intervening 27 years, it has become clear that this clinical-pathological correspondence is not always consistent. Extensive AD pathology, particularly diffuse amyloid plaques, can be present in the absence of any obvious symptoms [2–4]. Additionally, AD pathophysiology can manifest itself with clinically atypical presentations and prominent language and visuospatial disturbances [5–7]. Accordingly, in the revised NIA–Alzheimer's Association criteria, a semantic and conceptual distinction is made between AD pathophysiological processes (abbreviated as AD-P) and the various clinically observable syndromes that result (abbreviated as AD-C). Therefore, in this document and in the three associated diagnostic workgroup documents, a distinction is made between the syndromic labels that denote different qualitative and quantitative clinical expressions of disease (AD-C) and the pathophysiological process (AD-P) that underlies the syndrome.

Knowledge about the neuropathology of AD has also expanded over the past quarter century, and several sets of criteria for the neuropathological diagnosis of AD have been published, including an initial effort by the NIA [8], one from the Consortium to Establish a Registry for Alzheimer's Disease [9], and one from the NIA and Alzheimer's Association that is the one most widely used currently, the NIA–Reagan Institute criteria [10]. The current effort at redefining the clinical diagnosis of the preclinical and symptomatic disorders associated with AD-P assume that the fundamental characteristics of AD pathology—namely the presence of at least a moderate number of neuritic plaques containing β -amyloid in a low-power microscopic section of some region of neocortex and the extent of the regional distribution of neurofibrillary tangle pathology corresponding to Braak

and Braak stage IV or higher—will continue to define the neuropathological entity of AD.

Over the past 27 years, it has become abundantly clear that the cognitive deficits that accompany AD-P evolve gradually. A now voluminous literature on mild cognitive impairment (MCI) has arisen since the mid 1990s that documents the gradual impairments of cognitive function that precede the point where significant interference in daily activities occur [11]. The clinical dementia rating scale likewise reflects gradual development of clinical disease severity [12]. The 1984 criteria did not account for cognitive impairment that did not reach the threshold for dementia. The 1984 one-to-one clinical-pathological correspondence model did not account for the fact that AD-P develops slowly over many years, with dementia representing the end stage of many years of pathology accumulation in those patients who do become demented. Moreover, intermediate levels of AD pathological severity map onto clinical impairment, which is intermediate between normality and dementia [13–16]. Issues pertaining to intermediate clinical and pathological states and pertaining to clinical-pathological discordance are addressed in the revised criteria. Two recent publications by an international working group also emphasize the importance of identifying individuals in this intermediate stage of disease [17,18].

Because knowledge of the non-AD dementias was considerably more rudimentary in 1984, the 1984 criteria were vague in defining how distinctions between AD dementia and the major alternatives should be made. A prevalent notion, which is no longer widely held, was that reversible systemic disorders (e.g., thyroid disease, B12 deficiency, and others) were common mimics of AD dementia [19]. The concept of Lewy body disease did not exist [20,21]. That there was an entity of frontotemporal dementia because of something other than Pick's disease (which was thought to be extremely rare and not diagnosable in life) was not a consideration [22–24]. Although the concept of aphasia owing to neurodegenerative disease had been described only 2 years earlier [25], its pathological and neuropsychological features would begin to crystallize only 20 years later [26]. The fact that neuropsychiatric symptoms can be associated with AD was also not widely appreciated at the time, despite the fact that Alzheimer's celebrated first case had prominent delusions. The common coexistence of covert cerebrovascular disease, Lewy body disease, and AD pathology in elderly persons was not appreciated [27,28]. Thus, it has been only in the past decade that a better understanding of the distinctions and overlaps of the non-AD dementias with AD have begun to emerge. These concepts are embodied in the revised criteria.

Genetic discoveries in familial forms of early onset AD indicate that the initiating molecular events ultimately leading to both clinical and pathological AD begin with disordered beta-amyloid ($A\beta$) metabolism [29]. Recent data suggest that although familial AD can be characterized by overproduction of $A\beta_{42}$, late-onset sporadic AD may be

characterized by decreased clearance of $A\beta$ [30]. The major genetic risk factor for late onset AD is the $\epsilon 4$ allele of the apolipoprotein E gene, which is involved in $A\beta$ trafficking [31]. Therefore, the available genetic risk data overwhelmingly point to the $A\beta$ amyloid pathway as the initiating, or at least a very early pathophysiological event in the disease cascade [32].

Various features of AD pathology have been shown to relate to clinical symptoms differently. Clinical-autopsy correlation studies demonstrate a much tighter correlation between neurofibrillary pathology and cognitive impairment than between amyloid pathology and cognitive impairment [33,34]. However, the aspect of AD pathology that is most closely coupled with cognitive impairment is neurodegeneration, particularly synapse loss [35–37]. Approximately 30% of cognitively normal elderly subjects have some level of AD-P, and many of these individuals meet neuropathologic criteria for AD despite being free of apparent cognitive symptoms [3,4,37]. This 30% figure nearly perfectly matches the observed frequency of “amyloid positivity” in studies of cognitively normal subjects aged >65 years with positron emission tomography (PET) amyloid imaging and cerebrospinal fluid (CSF) assays [38–42], and roughly corresponds to the prevalence of AD dementia approximately a decade later [43]. These observations have been interpreted by some to imply an ordered sequence to the development of AD-P and its clinical consequences [44–47]. Rather than developing simultaneously, amyloid pathology and neurodegenerative pathology, in the form of neurofibrillary tangle formation and neuronal/synapse loss, may occur on different time scales [44]. $A\beta$ pathology is thought to develop first during the long preclinical phase, whereas the development of neurofibrillary pathology accelerates slightly before the appearance of the symptomatic phase of AD [48].

3. Biomarkers of AD

Two notable differences from the AD criteria published in 1984 [1] are incorporation of biomarkers of the underlying disease state and formalization of different stages of disease in the diagnostic criteria. Biomarkers of various features of AD-P have been developed and are being validated [49]. Biomarkers are parameters (physiological, biochemical, anatomic) that can be measured *in vivo* and that reflect specific features of disease-related pathophysiological processes. Although in the past the term biomarker was most often used in reference to fluid analytes, it has been used in all three workgroup documents to describe both fluid and imaging measures. A variety of biomarkers are discussed in the three workgroup documents. Some are discussed in terms of potential future applications, for example, resting state functional network connectivity [50]. However, only the five most widely studied biomarkers of AD based on the current literature are formally incorporated into the diagnostic criteria at this time.

The probabilistic framework for the incorporation of biomarkers was discussed extensively within each of the workgroups, and by the members of the workgroups charged with harmonizing the approach to biomarkers across the three documents. There was a consensus on several fundamental issues. First, it was important to tie the biomarkers as closely as possible to the pathological criteria for AD. Evidence suggests that together, the buildup of β -amyloid protein in plaques and tau deposition in neurofibrillary tangles is associated with neuronal injury. Second, it was agreed that the specificity of any biomarker for AD needed to be incorporated into the diagnostic schema. Evidence suggests that although both $A\beta$ deposition and elevated tau/phosphorylated tau are hallmarks of AD, alterations in these proteins are seen in other neurological disorders. Because elevations in $A\beta$ seem to be more specific than alterations in tau, it was decided to divide the biomarkers into two major categories: (1) *the biomarkers of $A\beta$ accumulation*, which are abnormal tracer retention on amyloid PET imaging and low CSF $A\beta_{42}$, and (2) *the biomarkers of neuronal degeneration or injury*, which are elevated CSF tau (both total and phosphorylated tau); decreased fluorodeoxyglucose uptake on PET in a specific topographic pattern involving temporoparietal cortex; and atrophy on structural magnetic resonance—again in a specific topographic pattern—involving medial, basal, and lateral temporal lobes and medial and lateral parietal cortices.

Underlying the deliberations of the groups was the recognition that the onset and progression of AD biomarkers likely follows an ordered temporal pattern. Biomarkers of $A\beta$ amyloid are indicative of initiating or upstream events which seem to be most dynamic (i.e., deviate most significantly from normal) before clinical symptoms. Biomarkers of neuronal injury and neuronal dysfunction are indicative of downstream pathophysiological processes which become dynamic later. Current evidence suggests that amyloid biomarkers may become abnormal anywhere from 10 to 20 years before noticeable clinical symptoms. Biomarkers of neurodegeneration become dynamic at a later point; some studies suggest that this may be shortly before clinical symptoms first appear. Progression of clinical symptoms closely parallels progressive worsening of neurodegenerative biomarkers [45–48].

Thus, biomarkers are used in the revised definitions of AD in all three disease phases, but the role of biomarkers differs somewhat in each of these stages. In the *preclinical phase*, biomarkers are used to establish the presence of AD-P in research subjects with no or very subtle overt symptoms. In both the MCI and AD dementia criteria, clinical diagnoses are paramount and biomarkers are complimentary. The core clinical diagnostic criteria for MCI and AD dementia are completely operational in a setting where no access to biomarkers exists. The approach to incorporation of biomarkers is more conservative in the diagnostic paradigm for symptomatic patients (MCI and AD dementia) than in preclinical research subjects. In the *symptomatic predemen-*

tia, *MCI*, *phase* biomarkers are used to establish the underlying etiology responsible for the clinical deficit. Biomarker severity, particularly neuronal injury biomarkers, also indicates the likelihood of imminent progression to AD dementia. In the *dementia phase*, biomarkers are used to increase or decrease, depending on the results, the level of certainty that AD-P underlies the dementia in an individual. The two major classes of biomarkers are treated equivalently in the MCI and dementia criteria. In contrast, they are ranked in a temporal hierarchy in the preclinical criteria, in that amyloid biomarkers become abnormal first and neuronal injury biomarkers become abnormal later. This temporal ordering notion is central to the staging proposed in the preclinical research criteria. The more conservative use for biomarkers in symptomatic subjects was felt to be a judicious approach pending more definitive outcomes research in this area.

4. Relevance to clinical practice

The recommendations of the three working groups differ in terms of current relevance to clinical practice. The core clinical criteria of the recommendations regarding AD dementia and MCI because of AD are intended to guide diagnosis in the clinical setting. However, the recommendations of the preclinical AD workgroup are intended purely for research purposes, and do not have any clinical utility at this time.

In addition, there was a broad consensus within the workgroups that much additional work needs to be done to validate the application of biomarkers as they are proposed in the workgroup documents. For example, additional biomarker comparison studies are needed, as is a more thorough validation with postmortem studies, and the use of combinations of biomarkers in studies has been limited. Extensive work on biomarker standardization is needed before widespread adoption of these recommendations at any stage of the disease. All biomarkers exist as continuous measures of AD-P. Therefore, biomarker standardization must also include gaining a broad consensus on how to obtain results that are interpretable as clearly normal, clearly abnormal, and perhaps intermediate. Moreover, obtaining standardized, reliable, and reproducible diagnostic caliber readouts of biomarker tests must be possible in any setting in which biomarkers are applied—research, clinical trials, or clinical care.

In summary, the new criteria for AD are presented in three documents, although the process is a continuous one with sometimes difficult-to-define boundaries between each discrete category. The evidence for preclinical AD is based almost entirely on AD biomarkers. Criteria for the earliest symptomatic manifestations, the MCI stage, represent a sharpening of previous efforts to define MCI. The MCI criteria also define an entity of MCI owing to AD-P, based on the conjunction of the clinical diagnosis and the presence of AD-P biomarkers. Finally, a revision of the 1984 criteria for dementia because of AD is provided.

The criteria for probable AD dementia expand the breadth of the 1984 criteria and include biomarker enhancements to the diagnosis of AD dementia. Ultimately, it is hoped that the scientific knowledge gained over the past quarter of a century, leading to the reconceptualization of “Alzheimer’s disease” proposed by the NIA-Alzheimer’s Association workgroup, will result in improved diagnosis and ultimately in effective disease-modifying therapy.

Acknowledgments

Clifford Jack serves as a consultant for Eli Lilly, Eisai, and Elan; is an investigator in clinical trials sponsored by Baxter and Pfizer Inc; and owns stock in Johnson and Johnson. Marilyn Albert serves as a consultant to Genentech and Eli Lilly and receives grants to her institution from GE Healthcare. David Knopman serves on a Data Safety Monitoring Board for Lilly Pharmaceuticals; is an investigator for clinical trials sponsored by Elan Pharmaceuticals, Forest Pharmaceuticals, and Baxter Healthcare; and is deputy editor of *Neurology* and receives compensation for editorial activities. Guy McKhann serves on a Data Safety Monitoring Board for Merck. Reisa Sperling has served as a site investigator and/or consultant to several companies developing imaging biomarkers and pharmacological treatments for early AD, including Avid, Bayer, Bristol-Myers-Squibb, Elan, Eisai, Janssen, Pfizer, and Wyeth. Maria Carrillo is an employee of the Alzheimer’s Association and reports no conflicts. Bill Thies is an employee of the Alzheimer’s Association and reports no conflicts. Creighton Phelps is an employee of the U.S. Government and reports no conflicts.

References

- [1] McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer’s disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer’s disease. *Neurology* 1984;34:939–44.
- [2] Davis DG, Schmitt FA, Wekstein DR, Markesbery WR. Alzheimer neuropathologic alterations in aged cognitively normal subjects. *J Neuropathol Exp Neurol* 1999;58:376–88.
- [3] Knopman DS, Parisi JE, Salviati A, Floriach-Robert M, Boeve BF, Ivnik RJ, et al. Neuropathology of cognitively normal elderly. *J Neuropathol Exp Neurol* 2003;62:1087–95.
- [4] Price JL, Morris JC. Tangles and plaques in nondemented aging and “preclinical” Alzheimer’s disease. *Ann Neurol* 1999;45:358–68.
- [5] Alladi S, Xuereb J, Bak T, Nestor P, Knibb J, Patterson K, et al. Focal cortical presentations of Alzheimer’s disease. *Brain* 2007;130(Pt 10):2636–45.
- [6] Rabinovici GD, Jagust WJ, Furst AJ, Ogar JM, Racine CA, Mormino EC, et al. Abeta amyloid and glucose metabolism in three variants of primary progressive aphasia. *Ann Neurol* 2008;64:388–401.
- [7] Tang-Wai DF, Graff-Radford NR, Boeve BF, Dickson DW, Parisi JE, Crook R, et al. Clinical, genetic, and neuropathologic characteristics of posterior cortical atrophy. *Neurology* 2004;63:1168–74.
- [8] Khachaturian ZS. Diagnosis of Alzheimer’s disease. *Arch Neurol* 1985;42:1097–105.
- [9] Mirra SS, Heyman A, McKeel D, Sumi SM, Crain BJ, Brownlee LM, et al. The Consortium to Establish a Registry for Alzheimer’s Disease (CERAD). Part II. Standardization of the neuropathologic assessment of Alzheimer’s disease. *Neurology* 1991;41:479–86.
- [10] Hyman BT. The neuropathological diagnosis of Alzheimer’s disease: clinical-pathological studies. *Neurobiol Aging* 1997;18(4 Suppl):S27–32.
- [11] Petersen RC, Doody R, Kurz A, Mohs RC, Morris JC, Rabins PV, et al. Current concepts in mild cognitive impairment. *Arch Neurol* 2001;58:1985–92.
- [12] Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;43:2412–4.
- [13] Markesbery WR, Schmitt FA, Kryscio RJ, Davis DG, Smith CD, Wekstein DR. Neuropathologic substrate of mild cognitive impairment. *Arch Neurol* 2006;63:38–46.
- [14] Bennett DA, Schneider JA, Bienias JL, Evans DA, Wilson RS. Mild cognitive impairment is related to Alzheimer disease pathology and cerebral infarctions. *Neurology* 2005;64:834–41.
- [15] Jicha GA, Parisi JE, Dickson DW, Johnson K, Cha R, Ivnik RJ, et al. Neuropathologic outcome of mild cognitive impairment following progression to clinical dementia. *Arch Neurol* 2006;63:674–81.
- [16] Petersen RC, Parisi JE, Dickson DW, Johnson KA, Knopman DS, Boeve BF, et al. Neuropathologic features of amnesic mild cognitive impairment. *Arch Neurol* 2006;63:665–72.
- [17] Dubois B, Feldman HH, Jacova C, Cummings JL, Dekosky ST, Barberger-Gateau P, et al. Revising the definition of Alzheimer’s disease: a new lexicon. *Lancet Neurol* 2010;9:1118–27.
- [18] Dubois B, Feldman HH, Jacova C, Dekosky ST, Barberger-Gateau P, Cummings J, et al. Research criteria for the diagnosis of Alzheimer’s disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol* 2007;6:734–46.
- [19] Clarfield AM. The decreasing prevalence of reversible dementias: an updated meta-analysis. *Arch Intern Med* 2003;163:2219–29.
- [20] McKeith IG, Dickson DW, Lowe J, Emre M, O’Brien JT, Feldman H, et al. Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology* 2005;65:1863–72.
- [21] McKeith IG, Galasko D, Kosaka K, Perry EK, Dickson DW, Hansen LA, et al. Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the consortium on DLB international workshop. *Neurology* 1996;47:1113–24.
- [22] Gustafson L. Clinical picture of frontal lobe degeneration of non-Alzheimer type. *Dementia* 1993;4:143–8.
- [23] Neary D. Non Alzheimer’s disease forms of cerebral atrophy. *J Neurol Neurosurg Psychiatry* 1990;53:929–31.
- [24] Neary D, Snowden JS, Gustafson L, Passant U, Stuss D, Black S, et al. Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 1998;51:1546–54.
- [25] Mesulam MM. Slowly progressive aphasia without generalized dementia. *Ann Neurol* 1982;11:592–8.
- [26] Gorno-Tempini ML, Hillis AE, Weintraub S, Kertesz A, Mendez M, Cappa SF, et al. Classification of primary progressive aphasia and its variants. *Neurology* 2011;76:1006–14.
- [27] Schneider JA, Arvanitakis Z, Bang W, Bennett DA. Mixed brain pathologies account for most dementia cases in community-dwelling older persons. *Neurology* 2007;69:2197–204.
- [28] White L. Brain lesions at autopsy in older Japanese-American men as related to cognitive impairment and dementia in the final years of life: a summary report from the Honolulu-Asia aging study. *J Alzheimers Dis* 2009;18:713–25.
- [29] Hardy JA, Higgins GA. Alzheimer’s disease: the amyloid cascade hypothesis. *Science* 1992;256:184–5.
- [30] Mawuenyega KG, Sigurdson W, Ovod V, Munsell L, Kasten T, Morris JC, et al. Decreased clearance of CNS beta-amyloid in Alzheimer’s disease. *Science* 2010;330:1774.
- [31] Roses AD. Apolipoprotein E affects the rate of Alzheimer disease expression: beta-amyloid burden is a secondary consequence dependent on APOE genotype and duration of disease. *J Neuropathol Exp Neurol* 1994;53:429–37.

- [32] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002; 297:353–6.
- [33] Gomez-Isla T, Hollister R, West H, Mui S, Growdon JH, Petersen RC, et al. Neuronal loss correlates with but exceeds neurofibrillary tangles in Alzheimer's disease. *Ann Neurol* 1997;41:17–24.
- [34] Bennett DA, Schneider JA, Wilson RS, Bienias JL, Arnold SE. Neurofibrillary tangles mediate the association of amyloid load with clinical Alzheimer disease and level of cognitive function. *Arch Neurol* 2004; 61:378–84.
- [35] DeKosky ST, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol* 1990;27:457–64.
- [36] Terry RD, Masliah E, Salmon DP, Butters N, DeTeresa R, Hill R, et al. Physical basis of cognitive alterations in Alzheimer's disease: synapse loss is the major correlate of cognitive impairment. *Ann Neurol* 1991; 30:572–80.
- [37] Savva GM, Wharton SB, Ince PG, Forster G, Matthews FE, Brayne C. Medical Research Council Cognitive Function and Ageing Study. Age, neuropathology, and dementia. *N Engl J Med* 2009;360:2302–9.
- [38] Mintun MA, Larossa GN, Sheline YI, Dence CS, Lee SY, Mach RH, et al. [¹¹C]PIB in a nondemented population: potential antecedent marker of Alzheimer disease. *Neurology* 2006;67:446–52.
- [39] Aizenstein HJ, Nebes RD, Saxton JA, Price JC, Mathis CA, Tsopelas ND, et al. Frequent amyloid deposition without significant cognitive impairment among the elderly. *Arch Neurol* 2008; 65:1509–17.
- [40] Jack CR Jr, Lowe VJ, Senjem ML, Weigand SD, Kemp BJ, Shiung MM, et al. 11C PiB and structural MRI provide complementary information in imaging of Alzheimer's disease and amnesic mild cognitive impairment. *Brain* 2008;131(Pt 3):665–80.
- [41] Fagan AM, Mintun MA, Mach RH, Lee SY, Dence CS, Shah AR, et al. Inverse relation between in vivo amyloid imaging load and cerebrospinal fluid Abeta42 in humans. *Ann Neurol* 2006; 59:512–9.
- [42] Shaw LM, Vanderstichele H, Knapik-Czajka M, Clark CM, Aisen PS, Petersen RC, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol* 2009;65:403–13.
- [43] Rowe CC, Ellis KA, Rimajova M, Bourgeat P, Pike KE, Jones G, et al. Amyloid imaging results from the Australian Imaging, Biomarkers and Lifestyle (AIBL) study of aging. *Neurobiol Aging* 2010; 31:1275–83.
- [44] Ingelsson M, Fukumoto H, Newell KL, Growdon JH, Hedley-Whyte ET, Frosch MP, et al. Early Abeta accumulation and progressive synaptic loss, gliosis, and tangle formation in AD brain. *Neurology* 2004;62:925–31.
- [45] Jack CR Jr, Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW, et al. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* 2010;9:119–28.
- [46] Mormino EC, Kluth JT, Madison CM, Rabinovici GD, Baker SL, Miller BL, et al. Episodic memory loss is related to hippocampal-mediated beta-amyloid deposition in elderly subjects. *Brain* 2009; 132(Pt 5):1310–23.
- [47] Perrin RJ, Fagan AM, Holtzman DM. Multimodal techniques for diagnosis and prognosis of Alzheimer's disease. *Nature* 2009;461:916–22.
- [48] Jack CR Jr, Lowe VJ, Weigand SD, Wiste HJ, Senjem ML, Knopman DS, et al. Serial PIB and MRI in normal, mild cognitive impairment and Alzheimer's disease: implications for sequence of pathological events in Alzheimer's disease. *Brain* 2009;132(Pt 5):1355–65.
- [49] Hampel H, Frank R, Broich K, Teipel SJ, Katz RG, Hardy J, et al. Biomarkers for Alzheimer's disease: academic, industry and regulatory perspectives. *Nat Rev Drug Discov* 2010;9:560–74.
- [50] Buckner RL, Snyder AZ, Shannon BJ, LaRossa G, Sachs R, Fotenos AF, et al. Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J Neurosci* 2005;25:7709–17.