

References:

- Chhabra, P.T., Rattinger, G.B., Dutcher, S.K., Hare, M.E., Parsons, K.L., Zuckerman, I.H.. (2012). Medication reconciliation during the transition to and from long-term care settings: A systematic review. *Research in Social and Administrative Pharmacy*, 8, 60-75.
- Dougherty, P., Katz, P. (2005). Complementary and alternative care in the long-term care setting. *Annals of Long Term Care*, 13. Available at www.annalsoflongtermcare.com/article/4003
- Fick, D., Semla, T., Beizer, J., Brandt, N., Dombrowski, R., DuBeau, C., et al., (2012). AGS updates Beers criteria for potentially inappropriate medication use in older adults. Available at: http://www.americangeriatrics.org/health_care_professionals/clinical_practice/clinical_guidelines_recommendations/2012
- Handler, S.M., Hanlon, J.T., Perera, S., Roumani, Y.F., Nace, D.A., Fridsma, D.B., Saul, M.I., Castle, N.G., and Studenski, S.A., (2008). Consensus list of signals to detect potential adverse drug reactions in nursing homes. *Journal of the American Geriatrics Society*, 56, 808-15.
- Institute of Medicine. Preventing medication errors. Available at: www.iom.edu/Reports/2006/Preventing-Medication-Errors-Quality-Chasm-Series.aspx
- MacKinnon, N. J., Kaiser, R. M., Griswold, P., and Bonner, A. (2009). Medication reconciliation and seamless care in the long term care setting. *Annals of Long Term Care*, 17, 36 – 40.
- Marcum, Z.A., Handler, S.M., Wright, R., and Hanlon, J.T. (2010). Interventions to improve suboptimal prescribing in nursing homes: A narrative review. *American Journal of Geriatric Pharmacotherapy*, 8, 183-200.
- Pham, C.B., and Dickman, R.L.. (2007). Minimizing adverse drug events in older patients. *American Family Physician*, 76, 1837-44.

