



## Temsirolimus (Torisel®)

**Pronounced:** TEM-sir-OH-li-mus

**Classification:** Kinase Inhibitor

### About Temsirolimus (Torisel®)

Temsirolimus is a kinase inhibitor which inhibits the mTOR kinase. A kinase is an enzyme that promotes cell growth. There are many types of kinases, which control different phases of cell growth. By blocking a particular enzyme from working, this medication can slow the growth of cancer cells.

### How to Take Temsirolimus

Temsirolimus is given intravenously (through an IV, into a vein). You may be given an antihistamine, such as diphenhydramine, prior to the infusion to decrease the chance of a reaction to the medication. Symptoms of a reaction include: flushing, facial swelling, shortness of breath, and chest pain. Let your nurse know right away if you experience any problems during the infusion.

The blood levels of this medication can be affected by certain foods and medications, so they should be avoided during treatment and between doses. These include: warfarin, dexamethasone, carbamazepine, rifampin, phenytoin, St. John's wort,azole antibiotics (fluconazole, itraconazole, ketoconazole, posaconazole, voriconazole), tacrolimus, grapefruit juice, phenobarbital and certain anti-retroviral therapies. Be sure to tell your healthcare provider about all medications and supplements you take.

You, or anyone you live with, should avoid having live or live-attenuated vaccines while receiving this medication. These include herpes zoster (Zostavax) for shingles prevention, oral polio, measles, nasal flu vaccine (FluMist®), rotovirus and yellow fever vaccines.

### Possible Side Effects of Temsirolimus

There are a number of things you can do to manage the side effects of temsirolimus. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### ***Low Red Blood Cell Count (Anemia)***

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

#### ***High Blood Sugar (Hyperglycemia)***

This medication can cause elevated blood sugar levels in patients with and without diabetes. Your oncology care team will monitor your blood sugar. If you develop increased thirst, urination or hunger, blurry vision, headaches or your breath smells like fruit, notify your healthcare team. Diabetics should monitor their blood sugar closely and report elevations to the healthcare team.

#### ***Infection Risk and Low White Blood Cell Count (Leukopenia or Neutropenia)***

This medication can suppress your immune system, putting you at higher risk of getting an infection. White blood cells (WBC)

are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4° F or 38° C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with, has any vaccinations.

### **Liver Toxicity**

This medication can cause liver toxicity, which your oncology care team may monitor for using blood tests called liver function tests. Notify your healthcare provider if you notice yellowing of the skin or eyes, your urine appears dark or brown, or you have pain in your abdomen, as these can be signs of liver toxicity.

### **Kidney Problems**

This medication can cause renal failure. If you notice that you are urinating less, or that your urine is dark in color, notify your healthcare provider. Your provider may also monitor the amount of protein in your urine while on this medication.

### **Fatigue**

[Fatigue](#) is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

### **Rash**

Some patients may develop a rash, scaly skin, or red itchy bumps. Use an alcohol free moisturizer on your skin and lips; avoid moisturizers with perfumes or scents. Your oncology care team can recommend a topical medication if itching is bothersome. If your skin does crack or bleed, be sure to keep the area clean to avoid infection. Be sure to notify your oncology care team of any rash that develops, as this can be a reaction. They can give you more tips on [caring for your skin](#).

### **Mouth Ulcers (Mucositis)**

Certain cancer treatments can cause [sores or soreness in your mouth](#) and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

### **Low Platelet Count (Thrombocytopenia)**

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib) etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding use of these agents and all over the counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

### ***Nausea and/or Vomiting***

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

### ***Peripheral Edema***

Peripheral edema is swelling of the extremities caused by retention of fluid. It can cause swelling of the hands, arms, legs, ankles and feet. The swelling can become uncomfortable. Notify your oncology care team if you are experiencing any new or worsening swelling.

### ***Decrease in Appetite and Taste Changes***

[Nutrition](#) is an important part of your care. Cancer treatment can affect your appetite and, in some cases, the side effects of treatment can make eating difficult. Ask your oncology care team about nutritional counseling services at your treatment center to help with food choices.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or find that food has no taste at all. You may dislike foods or beverages that you liked before receiving cancer treatment. These symptoms can last for several months or longer after treatment ends.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell. Sometimes cold food has less of an odor.
- Add extra flavor to meat or fish by marinating it in sweet juices, sweet and sour sauce or dressings. Use seasonings like basil, oregano or rosemary to add flavor. Bacon, ham and onion can add flavor to vegetables.

### ***Muscle or Joint Pain/Aches and Headache***

Your healthcare provider can recommend medications and other strategies to help relieve pain.

### ***Lung Changes***

This medication may cause a lung problem called interstitial lung disease. Notify your Healthcare provider if you have new or worsening shortness of breath, cough, wheezing or difficulty breathing.

### ***Diarrhea***

Your oncology care team can recommend medications to relieve [diarrhea](#). Also, try eating [low-fiber](#), bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole grain breads, cereals and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include: applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses of non-alcoholic, un-caffeinated fluid a day to prevent dehydration.

### ***Less common, but important side effects can include:***

- ***Increase in Cholesterol and Triglycerides:*** Patients receiving temsirolimus may develop increased cholesterol and triglyceride levels in the blood. Your provider will monitor for this. Some people may require a lipid lowering medication to treat this increase.

- **Wound Healing:** This medication can lead to slower or incomplete wound healing, such as a surgical wound not healing or staying closed. Therefore, it is recommended that the medication be used with caution before or after surgery. In addition, any surgical incision should be fully healed prior to starting or restarting the medication. If you have a surgical wound that has not healed or begins to have signs of infection (redness, swelling, warmth), report this to your healthcare team.
- **GI Tear:** This medication can cause a tear in the intestinal wall, also called a gastrointestinal perforation. Signs of this can include: new or worsening pain in the abdomen, new abdominal swelling, chills, fever, constipation, nausea or vomiting. If you experience any of these, contact your healthcare provider immediately or go to the emergency room.
- **Allergic Reactions:** In some cases, patients can have an allergic reaction to this medication. Signs of a reaction can include: shortness of breath or difficulty breathing, chest pain, rash, flushing or itching or a decrease in blood pressure. If you notice any changes in how you feel during the infusion, let your nurse know immediately. The infusion will be slowed or stopped if this occurs.

### **Reproductive Concerns**

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment and for at least 3 months after treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should not breastfeed while taking this medication or for 3 weeks after your last dose.

---

OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem or have questions or concerns about the medication that you have been prescribed, you should consult your health care provider.