



Etoposide (Toposar®, VePesid®, Etopophos®, VP-16)

Pronounced: e-TOE-poe-side

Classification: Topoisomerase inhibitor

About Etoposide (Toposar®, VePesid®, Etopophos®, VP-16)

Etoposide works by blocking the action of an enzyme in cells called topoisomerase, which is necessary for cell replication and tumor growth. Cells need this enzyme to keep their DNA in the proper shape when they are dividing. Blocking this enzyme leads to breaks in the DNA, which leads to cancer cell death.

How to Take Etoposide

Etoposide is given by intravenous (IV, into a vein) infusion. It can also be given by mouth in the form of a capsule. The dosage and schedule is determined by the person's size, type of cancer, and mode of administration. It can be given alone or with other medications.

Even when carefully and correctly administered by trained personnel, this drug may cause a feeling of burning and pain. There is a risk that this medication may leak out of the vein at the injection site, resulting in tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at any time during or after the injection, notify your care team immediately. Do not apply anything to the site unless instructed by your care team.

Possible Side Effects of Etoposide

There are a number of things you can do to manage the side effects of etoposide. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your **WBC count can drop**, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4° F or 38° C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- **Washing hands**, both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent **mouth care**.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with, has any vaccinations.

Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the **red cell count is low**, you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the **count is low** you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib) etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding use of these agents and all over the counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Mouth Ulcers (Mucositis)

Certain cancer treatments can cause **sores or soreness in your mouth** and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing **regular mouth care** can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage **nausea and vomiting**. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become **thin, brittle, or may fall out**. This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

Fatigue

Fatigue is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

Less common, but important side effects can include:

- **Lowering of Blood Pressure:** Blood pressure may drop while this medication is being infused. When receiving this medication through a vein, your nurse will be checking your blood pressure before and during the infusion. It may become necessary to stop the administration of this medication or slow down the infusion rate if your blood pressure drops.
- **Allergic Reactions:** Patients can have an allergic reaction to this medication. Signs of a reaction can include: swelling, chills, fever, increased heart rate, shortness of breath or difficulty breathing, or a decrease in blood pressure. If you notice

any changes in how you feel during the infusion, let your nurse know immediately. The infusion will be slowed or stopped if this occurs.

- **Radiation Recall:** Radiation recall is when the administration of a medication causes a skin reaction that looks like a sunburn (redness, swelling, soreness, peeling skin) in areas where radiation was previously given. Notify your oncology team if you notice this side effect. Treatment can include topical steroid ointments and a delay in your next chemotherapy dose.
- **Secondary Malignancies:** There is a very low risk of developing leukemia due to treatment with this medication, which can occur many years after treatment. This is most often associated with repeated treatments or high doses.

Sexual & Reproductive Concerns

This medication may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment and for at least 6 months for women and 4 months for men after treatment, even if your menstrual cycle stops or you believe you are not producing sperm. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. You should not breastfeed while receiving this medication.

OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem or have questions or concerns about the medication that you have been prescribed, you should consult your health care provider.