Ernestina Schipani, M.D., Ph.D. William Wikoff Smith Professor of Orthopaedic Surgery Full Professor of Orthopedic Surgery University of Pennylvania Perelman Medical School 3450 Hamilton Walk, 310A Stemmler Hall, Philadelphia, PA Work Phone: (215) 898-0045 Ernestina.Schipani@Pennmedicine.upenn.edu

Date of Birth	June 19, 1962
Place of Birth	Catanzaro, Calabria, Italy
Citizenship	Italian and USA

Education and Training

10/1/75-07/1/79	High School Diploma, <u>60/60 (highest score)</u> , Liceo Classico Pitagora,
	Crotone, Italy
10/1979-10/1985	M.D., <u>Summa Cum Laude</u> , Medical School of Pisa and Sant'Anna
	School of Advanced Studies, Pisa, Italy
11/1985-07/1988	Specialty Degree in Endocrinology, Summa Cum Laude, Medical
	School of Pisa, University of Pisa, Pisa, Italy
11/1985-12/1989	Ph.D., Summa Cum Laude, Sant'Anna School of Advanced Studies,
	Pisa, Italy
09/1987-10/1987	Visiting Fellow, Department of Internal Medicine, Bone Division
	Jewish Hospital, Washington University, St. Louis, MO
02/1990-10/1993	Research Fellow, Department of Internal Medicine, Massachusetts
	General Hospital-Harvard Medical School, Boston, MA
11/1992-03/1993	Visiting Fellow, Department of Molecular Pathophysiology, National
	Institutes of Health, Bethesda, MD
07/1994-07/1996	Research Fellow, Department of Internal Medicine, Massachusetts
	General Hospital, Boston, MA

Academic, Administrative, and Clinical Appointments

Academic Appointments

10/1993-11/1993	Instructor, Division of Endocrinology, Department of Internal
	Medicine, Harvard Medical School, Boston, MA
12/1993-05/1994	Assistant Professor of Medicine (with tenure), Division of
	Endocrinology, Department of Internal Medicine, University of Pisa
	Medical School of Pisa, Pisa, Italy
06/1994-06/1997	Instructor, Division of Endocrinology, Department of Internal
	Medicine, Massachusetts General Hospital -Harvard Medical School,
	Boston, MA
07/1997-05/2006	Assistant Professor of Medicine, Division of Endocrinology,
	Department of Internal Medicine, Massachusetts General Hospital -
	Harvard Medical School, Boston, MA

05/2006-06/2011	Associate Professor of Medicine, Division of Endocrinology, Department of Internal Medicine, Massachusetts General Hospital - Harvard Medical School Poston, MA
07/2011-08/2013	Professor of Medicine (with tenure) , Division of Endocrinology, Department of Internal Medicine, Indiana University Medical School, Indianapolis, IN
07/2011-08/2013	Professor of Anatomy and Cell Biology , Department of Anatomy and Cell Biology, Indiana University Medical School, Indianapolis, IN
09/2013-10/2020	Professor of Orthopaedic Surgery (with tenure) , Department of Orthopaedic Surgery, University of Michigan Medical School, Ann Arbor, MI
09/2013-10/2020	Professor of Medicine, Division of Endocrinology, Department of Medicine, University of Michigan Medical School, Ann Arbor, MI
03/2015-10/2020	Professor of Cell and Developmental Biology , University of Michigan Medical School, Ann Arbor, MI
11/2020-present	William Wikoff Professor of Orthopedic Surgery, Full Professor of Orthopedic Surgery (with tenure), University of Pennsylvania, Perelman School of Medicine, Philadelphia, PA

Other Appointments

Assistant in Biology, Department of Internal Medicine,
Massachusetts General Hospital, Boston, MA
Associate in Biology, Department of Internal Medicine,
Massachusetts General Hospital, Boston, MA
Consultant, Department of Internal Medicine, Massachusetts
General Hospital, Boston, MA
Full Member, Melvin and Bren Cancer Center, Indiana
University Medical School, Indianapolis, IN
Member, Center for Organogenesis, University of Michigan
Medical School, Ann Arbor, MI
Member, Institute for Regenerative Medicine, University of
Pennsylvania, Perelman School of Medicine, Philadelphia, PA

Licensure and Certifications

11/1985	Italian Medical License
12/1992	ECFMG

Research Interests

My laboratory focuses on skeletal development, leveraging insights from developmental biology to better understand skeletal diseases and identify potential new treatments.

Early in my career, I cloned the human PTH/PTHrP receptor and its gene, uncovering that gain-offunction mutations in this receptor are responsible for Jansen Metaphyseal Chondrodysplasia—a severe form of short-limbed dwarfism associated with hypercalcemia. Using mouse models carrying these mutations, I contributed to defining the PTH/PTHrP receptor's critical role in skeletal biology. Subsequently, I pioneered the concept that oxygen gradients regulate tissue morphogenesis during skeletal development. Beyond serving as a key metabolic substrate in enzymatic reactions such as mitochondrial respiration, oxygen also functions as a regulatory signal. My laboratory investigates how hypoxia and hypoxia-driven pathways influence skeletal development, aiming to uncover novel mechanisms of cellular adaptation to low oxygen levels and identify therapeutic targets for cartilage and bone diseases.

To achieve these goals, we employ genetically modified mouse models and a wide array of in vivo, ex vivo, and in vitro approaches to analyze phenotypes and elucidate the underlying biology.

The PTH/PTHrP receptor (PTHR1) in development and disease

I was a member of the team who cloned the cDNAs encoding the rat and opossum parathyroid hormone (PTH)/PTH related peptide (PTHrP) receptors (*also known as PTHR1s*), and I cloned the human homolog of this receptor and its gene. My study solved a long-lingering question in the field by proving that the PTH/PTHrP receptors expressed in bone and kidney are identical proteins. Next, I demonstrated that, at odds with what had been for a long time hypothesized, Pseudohypoparathyroidism type 1b, a rare endocrine disorder of calcium and phosphate homeostasis, was not caused by mutations in the PTH/PTHrP receptor gene. This finding prompted a wide genome search that eventually led to the identification of the Gsalpha gene as the one responsible for Pseudohypoparathyroidism type 1b.

More importantly, I discovered that gain-of-function mutations of the PTH/PTHrP receptor result in Jansen Metaphyseal Chondrodysplasia, a severe form of short-limbed dwarfism associated with hypercalcemia. Jansen Metaphyseal Chondrodysplasia has been one of the first examples in the literature of a human disease being caused by a constitutively active G-protein coupled receptor. Taking advantage of the mutations I had identified in patients, I generated transgenic mice expressing a constitutively active PTH/PTHrP receptor (Jansen receptor) in chondrocytes and osteoblasts, respectively. Lessons from these transgenic mice have contributed to shaping up our current understanding of the role of the PTH/PTHrP receptor in cartilage and bone development and homeostasis, and hematopoiesis.

My laboratory is currently collaborating with my former mentor Dr. Harald Jueppner at MGH-Harvard Medical School to identify potential therapeutic avenues for the treatment of Jansen Metaphyseal Chondrodysplasia (Awarded R01 DK113039, Dr. Schipani Co-Investigator, Dr. Jueppner PI).

The hypoxia signaling pathway in development ans disease

HIF-1 and the reprogramming of metabolism in endochondral bone development

While studying the fetal growth plate, we became intrigued by its avascular nature, which led us to uncover the critical role of hypoxia-signaling pathways in skeletal development. Oxygen, beyond being a vital metabolic substrate, also functions as a key regulatory signal. We were among the first to propose that oxygen gradients are essential for tissue morphogenesis during skeletal development. Through our research, we discovered that the murine fetal growth plate exhibits a gradient of oxygenation, with a hypoxic core region. To further investigate, we developed the first conditional knockout model of hypoxia-inducible factor-1 alpha (HIF-1), providing definitive evidence that HIF-1 acts as a survival factor for hypoxic chondrocytes in the growth plate in vivo. The role of HIF-1 as a survival factor has since been confirmed in a variety of settings, including cancer models. We also established that HIF-1 is crucial for the timely differentiation of mesenchymal cells into chondrocytes and for joint development in vivo. Furthermore, we provided genetic evidence that vascular endothelial growth factor (VEGF)-a classical downstream target of HIF-1 and a known survival factor for chondrocytes—plays only a modest role in mediating HIF-1's survival function in cartilage. Instead, we showed that HIF-1-dependent metabolic reprogramming is a critical downstream mechanism supporting chondrocyte survival and differentiation. Notably, we demonstrated that HIF-1 reduces mitochondrial respiration and oxygen consumption in growth plate chondrocytes, a key adaptation ensuring their survival and proper differentiation under hypoxic conditions. Currently, we are investigating how the interplay between oxidative phosphorylation and HIF-1-mediated metabolic reprogramming governs skeletal development.

HIF1 and the reprogramming of metabolism in somitogenesis

In collaboration with Dr. Mark Lewandoski at NCI, we recently established that HIF-1 is critical for spine development as its loss in the presomitic mesoderm impairs somitogenesis and causes spine and rib malformations that closely mimic those observed in patients with Jarcho-Levin Syndrome, a rare form of spondylothoracic dysplasia. A manuscript reporting those findings is in preparation.

We are currently investigating whether the impairment of somitogenesis secondary to loss of HIF-1 is due to dysregulation of glycolysis and mitochondrial function in the presomitic mesoderm.

HIF-2 and the control of bone mass accrual and homeostasis

A gradient of oxygenation exists within the bone marrow, despite its high vascularization. This phenomenon underscores the complexity of the bone marrow microenvironment, where localized hypoxia plays critical roles in cellular function. In collaboration with Dr. Amato Giaccia at Stanford, we demonstrated that osteoblasts have the remarkable ability to produce and secrete erythropoietin (EPO), a hormone essential for erythropoiesis, when hypoxia-inducible factor-2 (HIF-2), another transcription factor central to the hypoxic response, is activated in those cells. Our findings suggest that transient pharmacological activation of the hypoxia signaling pathway in osteoblasts could serve as a therapeutic approach to enhance EPO production, especially in conditions characterized by EPO deficiency, such as chronic kidney disease. In addition to its role in EPO production, HIF-2 has distinct effects on skeletal development. Unlike HIF-1, HIF-2 plays a less critical role in growth plate development. Interestingly, however, our research uncovered that reducing or inhibiting HIF-2 in mesenchymal progenitors and their descendants during development leads to the formation of stronger bones with increased trabecular bone mass. This improvement is driven by the expansion of mesenchymal progenitor cells within the bone marrow, which subsequently differentiate into osteoblasts. By increasing the pool of these progenitors, we effectively enhanced bone-forming capacity. A particularly exciting breakthrough involved the testing a HIF-2 inhibitor, PT2399, which was originally developed to treat clear cell renal carcinoma. In mouse models of menopause (characterized by estrogen deficiency), this drug prevented bone loss and promoted bone formation by expanding the pool of early osteoblast precursors. These findings highlight a promising therapeutic avenue for addressing osteoporosis and other bone loss conditions by targeting HIF-2. Currently, our research employs unbiased methodologies, including transcriptomic analyses, to elucidate the molecular mechanisms underlying the expansion of mesenchymal progenitor cells when HIF-2 activity is pharmacologically suppressed. These studies aim to identify novel pathways that can be leveraged to optimize bone health therapeutics.

HIFs in the pathogenesis of fibroblastic tumors of the soft tissue and cartilage regeneration

Our laboratory has demonstrated that continuous activation of the hypoxia signaling pathway in mesenchymal progenitor cells of the limb bud may have detrimental effects on skeletal development. Specifically, this persistent activation leads to aggressive fibrosis in synovial joints, the development of fibroblastic tumors in soft tissue, and dwarfism by disrupting both the proliferation and hypertrophic differentiation of growth plate chondrocytes. These findings underscore the critical role of tightly regulated hypoxia signaling in maintaining skeletal health and normal development. Notably, we also discovered that activation of the hypoxia signaling pathway in mesenchymal progenitors results in the formation of ectopic cartilage in the soft tissues surrounding the growth plate and promotes matrix accumulation in the developing growth plate. These findings suggest that increased activity of the hypoxia signaling pathway may be sufficient to drive cartilage formation under certain conditions. Taken together, if appropriately controlled, transient activation of hypoxia signaling could be harnessed to promote cartilage regeneration. By stimulating chondrogenesis while inhibiting hypertrophy, this strategy could enable cartilage repair in vitro and in vivo without inducing adverse outcomes, such as synovial fibrosis or fibroblastic tumor formation, provided the activation is carefully timed and transient. Our current research focuses on two main objectives: 1. Investigating the role of the hypoxia signaling pathway in the initiation and progression of fibroblastic tumors of the soft tissue; 2. Attempting to appropriately exploit the hypoxia signaling pathway for regenerating cartilage. These efforts aim to elucidate the dual role of hypoxia signaling in both pathology and regeneration, offering insights into novel therapeutic strategies for conditions involving fibroblastic tumors and cartilage repair.

Grants

PRESENT and ACTIVE:

R01 HD112003 (Schipani, PI) NIH/NICHD	01/04/23-31/03/28 \$220,000	2.4 calendar months
Role: PI	-	
Title: Hypoxia and Mitochondria in	n Spine Development and O	Congenital Scoliosis
The goal of this study is to advance pathways, and bioenergetic metabolic	e our understanding of how olism control somitogenesis	hypoxia,hypoxia-driven
R01 DK113039 (Jueppner, PI)	09/15/18-06/30/29	
NIH/NIDDK	\$12,249 (Schipani)	0.84 calendar months
Role: Co-Investigator		
Title: PTH Inverse Agonists as The	erapy for Jansens' Disease	

The goal of this study is the identification of therapeutic avenues for the treatment of Jansen Metaphsyseal Chondrodysplasia.

R01 AR075770 (Ma, PI)	09/23/20-01/31/26	
NIH/NIAMS	\$250,000	1.2 calendar months
Role: Co-Investigator		

Title: Regenerating Hyaline Cartilage Using Nanofibrous Hollow Microspheres and Synergizing TGF-beta and HIF

The goal of this study is to establish a novel approach to promote chondrogenesis and prevent chondrocyte hypertrophy.

04/01/25-04/31/30

PENDING:

R01 AR084536 (Schipani, PI)

3.0 calendar months

NIH/NIAMS Role: PI

Title: *Mitochondrial Respiration and The Biology of Growth Plate Chondrocytes* The goal of this study is to establish how oxidative phosphorylation controls chondrocyte hypertrophy in the murine developing growth plate.

Major Previous Grants

1993-2000	PTH/PTHr Receptor Defects in Pseudohypoparathyroidism. RO1-DK4718 NIH/NIDDK Co-investigator
1996-2000	Constitutively active PTH/PTHrP receptors in vivo. RO1-DK5070 NIH/NIDDK Co-investigator

1997-2005	Parathyroid hormone and Osteoporosis: Therapy and Basic Mechanisms.P50-AR4485NIH/NIAMSPI Project VThe major goal: To use transgenic animals expressing a constitutively active PTH/PTHrPreceptor in cells of the osteoblast lineage to define how activation of this receptormodulates bone remodeling.
1997-2005	Parathyroid hormone and Osteoporosis: Therapy and Basic Mechanisms.P50-AR4485NIH/NIAMSPI Bone Analysis CoreThe major goal: to serve each of the individual projects of the SCOR in performing routine histology, in situ hybridization, immunohistochemistry, and hormonal measurements.
1999-2011	Developmental Regulation of Bone Morphogenesis. PO1- DK56246 NIH/NIDDK PI High Resolution Histology Core The major goal: to serve each of the individual projects in performing routine histology, in situ hybridization and immunohistochemistry.
2002-2006	Carboxyl-terminal PTH Receptors in Bone Cells. RO1-AR4706 NIH/NIAMS Co-investigator
2003-2011	Hormonal Control of Calcium Metabolism. PO1-DK11794 NIH/NIDDK Co-investigator Project V
2003-2011	Hormonal Control of Calcium Metabolism. PO1-DK11794 NIH/NIDDK Co-investigator, Tissue Phenotyping Core
2004-2006	PTH1R regulation in bone biology using mouse models. RO1-DK0228 NIH/NIDDK Co-investigator
2004-2005	Modulation of bone remodeling by administration of OPG, Alendronate or Zoledronate. Amgen PI The major goal: to investigate how treatment with OPG, alendronate or zoledronate affects bone remodeling in a mouse model of high bone turnover secondary to expression of a constitutively active PTH/PTHrP receptor in cells of the osteoblast lineage.
2005-2010	Specialized Center for Cell Based Therapy [SCCT]. U54-HL081030-01 PI Bone Analysis Core The major goal: to serve each of the individual projects in performing routine histology, in situ hybridization and immunohistochemistry
2008-2010	Modulation of bone remodeling by concomitant administration of OPG and PTH. Amgen PI

	The major goal: to study how administration of OPG modulates the bone marrow fibrosis in mice expressing a constitutively active PTH/PTHrP receptor in cells of the osteoblast lineage.
2011-2013	<i>Identification of a novel bone marrow population.</i> R21 AR060689 NIH/NIAMS PI The major goal: to study a novel bone marrow population recently identified in the bone marrow and its contribution to hone marrow fibrosis
2003-2014	Role of Hypoxia in Differentiation. RO1 AR058654 NIH/NIAMS PI The major goal: to study the role of hypoxia signaling pathways in endochondral bone development.
2009-2014	Notch and hypoxia in intervertebral disc development.RO1NIH/NIAMSCo-investigatorThe major goal: to study the role of Notch and hypoxia signaling pathways in intervertebral disc development.
2013-2018	Role of HIF-1 in intervertebral disc function RO1 AR055655 NIH/NIAMS Co-investigator The major goal of this project is to investigate the role of HIF-1 in intervertebral disc development and function.
2015-2017	 Exploring the physiological role of osteoblastic Epo and osteoblastic EpoR R21 AR067330 NIH/NIAMS PI The major goal: to study the role of osteoblastic EPO and osteoblastic EPOR in bone and bone marrow development and homeostasis.
2016-2017	Suppression of b-catenin hyperactivity by HIFs: Implications in colon cancer therapeutics. McCubed University of Michigan Co-PI The major goal: to provide critical mouse models for the execution of the study.
2016-2017	Role of the Hypoxia Signaling Pathway in Spine Development. Pediatric Orthopaedic Society of North America Co-investigator The major goal: to study whether loss of HIF-1a in somites alters somitogenesis and results in spine deformities, and eventually scoliosis.
2013-2019	<i>HIF-1alpha, a survival and differentiation factor for cartilage.</i> RO1 AR065403 NIH/NIAMS PI

The goal of this project is to study the role of mitochondrial metabolism downstream of HIF1 as a survival and differentiation factor for chondrocytes.

2016-2020	 P30 Michigan Integrative Musculoskeletal Health Core Center NIH/NIAMS Histology Core Director The Michigan Integrative Musculoskeletal Health Core Center (MiMHC) will enable vertically integrative, multi-scale musculoskeletal scince by increasing access to critical, specialized resources and expertise that are fundamental to the musculoskeletal research programs of center investigators.
2022	2022 Bone and Teeth Gordon Research Conference and Seminar R13 AR080434 NIH/NIAMS PI Major Goals: The skeleton is an organ with numerous vital functions. The 2022 Bones anD Teeth Gordon Research Conference and Seminar will discuss major new discoveries regarding the skeleton and its functions. Those new and exciting discoveries may lead to novel treatments of devastating diseases including osteoporosis, bone cancer, genetic disorders of the skeleton, and skeletal problems associated to diabetes.
2018-2023	 HIF-2alpha, a Novel Regulator of Osteoblastogenesis R01 AR073022 NIH/NIAMS PI The goal of this study is to determine the role of HIF2 in the regulation of bone mass and osteoblastogenesis.
2019-2024	Mitochondria and TFAM in osteoblast biology R01 AR074079 NIH/NIAMS PI

The goal of this study is to determine the role of TFAM in osteoblast biology.

Honors and Awards

1985	M.D., Summa Cum Laude, Sant'Anna School of Advanced Studies and Medical School of Pisa, Pisa, Italy
1989	Ph.D., Summa Cum Laude, Sant'Anna School of Advanced Studies, Pisa, Italy
1990-1993	Fellowship, Ministero della Ricerca Scientifica, Italy. Special grant for a
	scientific studies abroad; awarded by the Italian government
1993	Concorso for Assistant Professor Position at University of Pisa, prestigious
	national competition for university faculty positions, Italian government
1994-1996	National Osteoporosis Foundation Fellowship
1995	Travel Award of the International Conference on Calcium Regulating
	Hormones
1995	Young Investigator Award, American Society for Bone and Mineral Research
2019	Paula Stern Achievement Award-ASBMR Esteemed Award
2019	Fellow of the ASBMR

2020	Co-Chair of 2020 Bone and Teeth Gordon Research Conferences
2022	Chair of 2022 Bone and Teeth Gordon Research Conferences

Memberships in Professional Societies

1992-present	American Society for Bone and Mineral Research
1	2000-2018, 2022, 2023 – Abstract reviewer
	1996, 1997, 2000, 2005, 2006, 2012, 2013, 2014, 2019, 2020, 2023 – Moderator
	ASBMR meeting
	2008 – Co-Chair State-of-the-Art Lecture A: Role of oxygen sensing pathways
	2014 – Chair Skeletal Development abstract review category
	2015 – Chair Skeletal Development abstract review category
	2015 – Co-chair Symposium "Metabolism of Bone Cells"
	2015 – Discussion Leader "Grant writing workshop"
	2015-2017 – Member Education and Membership Committee
	2015-2016 – Member Organizing Committee ASBMR 2016
	2017-2019 - Council Member
	2017 – Council Liaison Women Committee
	2017 – Member Annual Meeting Program Advisory Committee
	2018 – Council Liaison Development Committee
	2018 – Member Nominating Committee
	2018-2019 – Council Liaison Publications Committee
	2019 – Co-chair Symposium "Cutting Edge Concepts: CRISPR beyond the
	mouse"
	2019-2021 – Member Annual Meeting Advisory Committee
	2021-2023 – Member Publications Committee
	2022 – Member Award Committee
	2024- Chair Premeeting Symposium on "Energy Metabolism in Skeletal
	Development and Disease"
	2024-Member Advisory Committee
1994-2011	American Society for Endocrinology
1999	Moderator of one session Endocrine Society meeting
2002-2011	American Society of Matrix Biology
2004-present	Association of Osteobiology
2005-present	ASCI
2005-2011	The New York Academy of Sciences
	2007 – Moderator of session, 2 nd conference on Skeletal Biology and Medicine,
	The New York Academy of Sciences
	2011 – Moderator of session, 4 th conference on Skeletal Biology and Medicine,
	The New York Academy of Sciences
2007-2015	International Bone and Mineral Society

Editorial Positions, Boards, and Peer-Review Services

Grant Review Activities

1999-2000	Medical Research Council (MRC, Canada), Ad hoc reviewer
2006	VA Grants, Ad hoc reviewer
2006	Committee for Review of Research Proposals, Harvard Stem Cell Institute,
	Harvard Medical School, Ad hoc reviewer

2007	Pilot and feasibility grants, Yale Core Center for Musculoskeletal Disorders, Medical School, Yale University, Ad hoc reviewer
2007-2011	NIH SBSR Study Section Regular Member
2008	Pilot and Feasibility grants Center for Metabolic Bone Diseases University of
2000	Alabama Ad hoc reviewer
2008-2014	ASBMR Career Enhancement Award Committee
2008-2014	NIH NCL Call Biology Special Emphasis Ad has reviewer
2009	NILL MTE Study Special Emphasis, Ad hoc reviewer
2013	NIH-MITE Study Section, Ad noc reviewer
2013	NIH-Special Emphasis Panel" Musculoskeletal Development, Injury and
0014	Regeneration", Chair
2014	NIH-Special Emphasis Panel "ZRG1 MOSS-U03", Chair
2014-2018	NIH-MTE Study Section, Regular Member
2014	NIH-NIAMS, Roundtable discussion on the role of disc degeneration in back
	pain
2015-present:	Fibrodysplasia Ossificans Progressiva (FOP) Foundation, Ad hoc Reviewer
2015	Pilot Projects, Yale Diabetes Center, Medical School, Yale University, Ad hoc
	reviewer
2015	NIA- ZAG1 ZIJ-8 (M2) (PO1), Ad hoc reviewer
2015	NIAMS-ZRG1-MOSS-U02, Ad hoc reviewer
2017	NIAMS-Special Emphasys Panel. Ad hoc reviewer
2017	UMHS-PUHSC Joint Institute, University of Michigan, Ad hoc reviewer
2018-2022	NIH-SBDD Study Section Ad hoc reviewer
2018	NIAMS Special Emphasys Panels. Ad hoc reviewer
2018	Swiss National Science Foundation Ad hoc reviewer
2018-present	NIA PO1s Ad hoc reviewer
2010 present. 2019	Pilot and feasibility grants Department of Medicine Medical School
2017	Wahington University Ad hoc reviewer
2019	Pilot and Feasibility grants Dental School University of Michigan Ad hoc
2017	reviewer
2010	Pilot and fassibility grants Musculoskalatal Center Medical School University
2019	of Michigan Ad has reviewer
2010	Dilat and fassibility grants. Contar for Organogenesis. Medical School
2019	Luiversity of Michigan Adhaa reviewer
2021	NIDDY Deard of Scientific Courselars Deview
2021	NILDK Board of Scientific Coursefors Review
2021	NIH Directors Early Independence Awards, Ad Hoc Reviewer
2023	NIAMS-Special Emphasis Panel, R13/U13, Ad hoc Reviewer
2023	NIH- Special Emphasis Panel "ZRGT MSOS V (02) M", Chair
2023-2024	NIH-U19, Chair
2023	PhD Thesis Committee-University of KU Leuwen, Belgium, External Reviewer
2023	PhD Thesis Committee-University of Monash, Melbourne, Australia, External Reviewer
2023	Pilot Grant Musculoskeletal Center-Washington University. St.Louis. External
	Reviewer
2024	GWIS National Fellowhip Program, Ad hoc Reviewer

<u>Editorial Board</u>

01/2001-01/2016	BoneKEy (Nature Publishing Group)
01/2002-12/2005	Endocrinology
07/2004-05/2015	Journal of Bone and Mineral Research
01/2007-01/2021	Bone
01/2016-12/2020	Endocrinology
10/2021-present	Journal of Bone and Mineral Research

Editor

2014-2015	Section Editor-Current Osteoporosis Reports
2015	Scientific Reports (Nature Publishing Group)
2019-2020	Guest Editor-Bone
2021-2022	Section Editor-Current Osteoporosis Reports
2021-2023	Guest Editor-Bone Reports
2021-2023	Reviewing Editor-eLife
2021-2024	Associate Editor-JCI Insight
2024	Associate Editor-Endocrinology
2024-	Consulting Editor-JCI Insight

Ad-hoc manuscript reviewer

American Journal of Endocrinology and Metabolism Arthritis and Rheumatism Blood Bone Calcified Tissue International Cancer Research Development Developmental Biology Developmental Cell Cell Metabolism eLife EMBO FASEB Endocrinology Genes and Development Human Molecular Genetics **JBMR** Journal of Cell Biology Journal of Clinical Endocrinology and Metabolism Journal of Clinical Investigation Journal of Experimental Research Journal of Orthopedic Research Journal of Biological Chemistry Molecular Cancer Research MCB Molecular Endocrinology Nature Medicine Nature Communication Osteoarthritis and Cartilage **PLoS** Genetics PLoSone PNAS Science Science Signaling Stem Cells Science Translational Medicine Stem Cell Reports

Teaching

Teaching of Students in Courses

2000-2008	Tutor – Course of Pathophysiology in the Renal/Musculoskeletal/Endocrine Block 2 nd year Medical Students, Harvard Medical School, Boston, MA
2004-2005	Lecturer – Molecular mechanisms of bone and joint formation, 2 nd year Medical Students, Harvard Medical School, Boston, MA
2015	Lecturer – Molecular and Cellular Mechanisms of Cartilage and Bone Development, Clinical Residents-Department of Orthopaedic Surgery,
2015	Lecturer – Hypoxia signaling pathways in development, CDB 582/583 Course "Stem cells to regenerative biology", University of Michigan, Ann Arbor, MI
2016	Lecturer – Endocrine Regulation of Bone and Calcium/Phosphate homeostasis I, Clinical Residents-Department of Orthopaedic Surgery, University of
2016	Michigan, Ann Arbor, MI Lecturer – Endocrine Regulation of Bone and Calcium/Phosphate homeostasis II, Clinical Residents-Department of Orthopedic Surgery, University of Michigan, Ann Arbor, MI
2016	Lecturer – Spine Development, Clinical Residents-Department of Orthopaedic Surgery University of Michigan Ann Arbor MI
2016-2017	Lecturer – Permanent Cartilage and Bone, Musculoskeletal Course-Medical School University of Michigan Ann Arbor MI
2016-2017	Lecturer – Bone formation, Musculoskeletal Course-Medical School, University of Michigan, Ann Arbor, MI
2016-2019	Lecturer- Bone Formation, Musculoskeletal Course-Dental School, University of Michigan, Ann Arbor, MI
2016-present	Lecturer– Permanent Cartilage and Bone, Musculoskeletal Course-Dental School, University of Michigan, Ann Arbor, MI
2016-2017	Lecturer – Permanent Cartilage and Bone, Histology Course for graduate and undergraduate students, University of Michigan, Ann Arbor, MI
2016-2017	Lecturer-Histology Course for graduate and undergraduate students "Bone formation" University of Michigan Ann Arbor MI
2017	Lecturer- VitD and Rickets, Clinical Residents-Department of Orthopaedic Surgery University of Michigan Ann Arbor MI
2017-2019	Lecturer- Skeletal Development, Clinical Residents-Department of Orthopaedic Surgery University of Michigan Ann Arbor MI
2017-2019	Lecturer- Permanent Cartilage and Adult Bone, IInd year Medical Students, University of Michigan, Ann Arbor, MI

Laboratory and Other Research Supervisory and Training Responsibilities

Advisory role for 2-3 postdoctoral fellows
PI at the Histology Core, Endocrine Unit, MGH, Boston, MA
External advisor member of the PhD thesis defense committee-Tissue
Engineering Department, Stony Brook University, NY
Member of the PhD thesis defense committee, Harvard Dental School, MA
Member of the PhD qualifying exam committee, Department of Physiology,
Medical School, University of Michigan, Ann Arbor, MI
Member of the PhD qualifying exam committee, Department of Physiology, Medical School, University of Michigan, Ann Arbor, MI

2016-present	PI at the Histology Core, Orthopaedic Research Laboratory, Medical School,
-	University of Michigan, Ann Arbor, MI
2017	Mentor of a thesis with honors at College of Literature, Science and the Arts,
	University of Michigan, Ann Arbor, MI
2018	Member of a PhD thesis defense committee, Department of Orthopedic Surgery,
	Medical School, University of Michigan, Ann Arbor, MI
2018	Member of a PhD thesis defense committee, Department of Cell and
	Developmental Biology, University of Hong Kong, Hong Kong, China
2019	Member of a Master thesis defense committee, Department of Orthodontics,
	Dental School, University of Michigan, Ann Arbor, MI

Formally Supervised Trainees

2000	Anna Giovannetti, PhD/ Research Associate, University of Pisa, Pisa, Italy
2001-2002	Anja Maier, MD/ Private Practice Physician
1999-2002	Laura Calvi, MD/ Young Investigator Award-ASBMR, Haddad Investigator
	Award, K08 Award (NIH), Professor, Medical School, University of Rochester,
	NY
2003-2004	Riccardo Chiusaroli, PhD/ Head of Histology and Pathology, Rottapharm,
	Monza, Italy
2005-2008	Masanobu Ohishi, MD-PhD/ Assistant Professor with tenure, Medical School,
	University of Kiushu, Japan
2005-2006	Sylvain Provot, PhD/Young Investigator Award-ASBMR, Assistant Professor
	with tenure, INSERM, Paris, France
2008	Ellinoora Aro, MD-PhD/Resident General Surgery, University of Helsinki,
	Finland
2008-2011	Richa Khatri, MD/ Resident General Surgery, Western Michigan University,
	Kalamazoo, MI
2009-2010	Wanida Ono, DDS-PhD/Associate Professor, Dental School, University of
	Texas, Houston, TX
2009-2010	Elisa Araldi, PhD/ Assistant Professor, ETH Zurich, Zurich, Switzerland
2010-2011	Erinn Rankin, PhD/ Assistant Professor, Radiation Oncology, Stanford, CA
2011-2015	Laura Mangiavini, MD/Associate Professor, University of San Raffaele,
	Milano, Italy
2015-2017	Kavitha Ranganathan, MD/Young Investigator Award-ASBMR, Resident
	General Surgery, University of Michigan Medical School, Ann Arbor, MI
2015-2018	Angela Yao, PhD/Young Investigator Award-ASBMR, Assistant Professor,
	University of Science and Technology, Shenzen, China
2012-2020	Christophe Merceron, PhD/ Young Investigator Award-ASBMR Senior
	Medical Information Associate, Prime Vigilance, Ann Arbor, MI
2021-2022	Lorenzo Arboit, MD, PhD Student University of Strasbourg, Strasbourg, France
2017-2023	Mohammed Parvez-Khan, PhD/Travel Award-ASBMR, Research Assistant
2021 2024	Professor, University of Toledo, Toledo, OH
2021-2024	Elena Sabini. MD. PhD student University of Pisa, Pisa, Italy
2022-2024	Giulia Lanzolla, MD/Travel Award-ASBMR, Assistant Professor, University
	of Cagliari, Italy

Local and Regional Presentations and Lectures

1993	The human PTH/PTH receptor from cloning to human diseases, Endocrine
	Division Lectures Massachusetts General Hospital, Boston, MA
1993	The PTH/PTHrP receptor in Pseudohypoparathyrioidism, Pediatric Department
	Lectures Massachusetts General Hospital, Boston, MA
	1

1995	The PTH/PTHrP receptor in Jansen Metaphyseal Chondrodysplasia, Pediatric Endocrine Grand Rounds Massachusetts General Hospital, Boston, MA
2004	Jansen's PTH/PTHrP receptors in bone growth and remodeling, Endocrine Grand Rounds Massachusetts General Hospital, Boston, MA
2004	Jansen's PTH/PTHrP receptors in bone growth and remodeling, Endocrine Grand Rounds Brigham and Women's Hospital, Boston, MA
2004	Hypoxia, HIF-1 alpha and VHL in endochondral bone development, Grand Round-Harvard Dental School, Boston, MA
2005	Hypoxia and HIF-1 alpha in growth plate development, Orthopaedic Research Seminar Series 2004-2005 Children's Hospital Boston MA
2006	Constitutively active PTH/PTHrP receptors in bone stromal cells, 4th Symposium on Membrane Biology Massachusetts General Hospital Boston MA
2012	Hypoxia Signaling Pathway in Development and in Differentiation, Tumor Microenvironment Seminar Series-Cancer Center-IU School of Medicine, Indianapolis, IN
2012	HIFs and VHL in cartilage, bone and hematopoiesis, Bone Seminar Series, IU School of Medicine, Indianapolis, IN
2013	Hypoxia signaling pathways in organogenesis. MSK Seminar Series, October 9, University of Michigan, Ann Arbor, MI (Seminar)
2013	Hypoxia signaling pathways in cartilage, bone and hematopoiesis: an update. Endocrine Seminar Series, November 1, University of Michigan, Ann Arbor, MI (Seminar).
2014	HIFs and VHL in cartilage and bone development. Orthopaedic Surgery Grand Rounds, February 11, University of Michigan, Ann Arbor, MI (Seminar)
2014	Hypoxia signaling pathways in development and differentiation. Seminar Series, December 9, Nephrology Division, University of Michigan, Ann Arbor, MI (Seminar)
2015	Hypoxia signaling pathways in development and differentiation. Seminar Series, January 15, Division of Hematology Oncology, University of Michigan, Ann Arbor, MI (Seminar)
2015	Hypoxia signaling pathways in development of the nucleus pulposus. April 27, The Frederick J. Fisher Pediatric Orthopaedic Lectureship, University of Michigan, Ann Arbor, MI (Invited Talk)
2016	Role of HIF-1a in spine development. April 25, The Frederick J. Fisher Pediatric Orthopaedic Lectureship, University of Michigan, Ann Arbor, MI (Invited Talk)
2016	HIFs and the osteogenesis-angiogenesis coupling. May 6, T32 Training Grant Seminar Series, Plastic Surgery University of Michigan, Ann Arbor, MI (Seminar)
2017	Impairment of mitochondrial respiration promotes survival of hypoxic chondrocytes. March 3rd, Endocrine Grand Round, Department of Medicine- Endocrinology, School of Medicine, University of Michigan, Ann Arbor, MI (Seminar)
2017	Ìmpairment of mitochondrial respiration promotes survival of hypoxic chondrocytes. "Brown Bag" Seminar Series, Department of Cell and Developmental Biology, School of Medicine, University of Michigan, Ann Arbor, MI (Seminar)
2107	survival of hypoxic cells <i>in vivo</i> . Department of Physiology School of Medicine, University of Michigan, Ann Arbor, MI (Seminar)
2018	Enhancing mitochondrial respiration causes severe intracellular hypoxia and death of hypoxic cells. Brown Bag" Seminar Series, Department of Cell and

2010	Developmental Biology, School of Medicine, University of Michigan, Ann
2019	Hypoxia, HIFs and mitochondria in skeletal development. Brown Bag" Seminar
	Series, Department of Cell and Developmental Biology, School of Medicine,
2021	University of Michigan, Ann Arbor, Mi (Seminar)
	Seminar, Perelman School of Medicine, University of Pennsylvania, Philadelphia,
	PA (Seminar)
2022	How hypoxia and Mitochondria shape up the skeleton. PCMD, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA (Short talk, Invited)
2023	How hypoxia and Mitochondria shape up the skeleton. IRM Annual Retreat, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA
	(Short Talk, invited)
2023	How hypoxia and Mitochondria shape up the skeleton. IDOM Seminar, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA (Seminar)

Committee, Organizational, and Volunteer Services

Local

2004-2005	ECOR Subcommittee for Review of Research Proposals, Massachusetts
	General Hospital, Boston, MA
2000-2010	SRAC, Massachusetts General Hospital, Boston, MA
2014-2015	RO1 Boot Camp, UMICH, Ann Arbor, MI
2014-present:	RAC Committee, Department of Orthopedics, Medical School, University of Michigan, Ann Arbor, MI
2014-2015	LAUNCH Committee, Dental School, University of Michigan, Ann Arbor, MI
2015-present:	BioArtography Committee, Department of Cell and Developmental Biology, Medical School, University of Michigan, Ann Arbor, MI
2015-2016	Award Nominating Committee, Department of Cell and Developmental Biology, Medical School, University of Michigan, Ann Arbor, MI
2015-2016	Graduate Students Admissions Committee, Department of Cell and Developmental Biology, Medical School, University of Michigan, Ann Arbor, MI
2016-2017	LAUNCH Committee, Dental School, University of Michigan, Ann Arbor, MI
2017-2108	Search Committee for Faculty Positions, Dental School, University of Michigan, Ann Arbor, MI
2018	Committee for abstract selection, Annual Symposium, Department of - Medicine:University of Michigan, Ann Arbor, MI
2019-2020	Search Committee for Faculty Positions, Department of Cell and Developmental Biology, Medical School, University of Michigan, Ann Arbor, MI
2019-2020	Search Committee for Faculty Positions, Dental School, University of Michigan, Ann Arbor, MI
2024-	Chair DCOAP Committee, Department of Orthopedic Surgery, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA

National and International Committees

1994 Scientific Committee for the 4th International Symposium on Endocrinology under 35

2007-2011	Scientific Committee for IBMS Davos, Workshop: Bone Biology &
06/07-06/2011	Interapeutics, International Bone of Mineral Society Member IBMS Council International Bone Mineral Society
2009-2011	Scientific Committee IBMS meeting 2011, International Bone of Mineral
2007 2011	Society
2009-2010	Scientific Committee Parathyroids 2010
03/2011	Discussion Leader, Gordon Conference on Cartilage Biology and Pathology,
	March 6-11, 201, Ventura, CA
06/2012-06/2015	Member Nominating Committee, International Bone Mineral Society
04/2013	Discussion Leader, Gordon Conference on Cartilage Biology and Pathology,
	April 7-11, 2013, Les Diablerets, Switzerland
09/2014-present:	Board Member of the Osteobiology Association
03/2015	Discussion Leader, Gordon Conference on Cartilage Biology and Pathology,
	March 22-27, 2015, Galveston, TX
2015-2016	ASBMR Organizing Committee
2015-2017	ASBMR Education and Membership Committee
2016-2017	ASBMR Annual Meeting Program Advisory Committee
2017-2019	ASBMR Council
2017	ASBMR Council Liason Women Committee
2018	ASBMR Council Liason Development Committee
2018	ASBMR Nominating Committee
2019	ASBMR Council Liason Publications Committee
2019-2020	ASBMR Annual Meeting Advisory Committee
2021-2023	ASBMR Publication Committee
2024	ASBMR Advisory Committee

Visiting Professorships, Seminars, and Extramural Invited Presentations

<u>National</u>

1992	The human PTH/PTH receptor: from cloning to human diseases.
	Molecular Pathophysiology Branch, NIH, Bethesda, MD (Seminar)
1999	Constitutively active PTH/PTHrP receptors: in vitro studies.
	ASBMR, Workshop on structure function of the PTH/PTHrP receptor St. Louis,
	MO (Seminar)
1999	The PTH/PTHrP receptor in endochondral bone development.
	American Society of Nephrology, Miami, FL (Lecture)
2000	Jansen PTH/PTHrP receptor in bone development.
	Endocrine Grand Rounds, NIH, Bethesda, MD (Visiting Professor)
2000	The PTH/PTHrP receptor in endochondral bone development.
	American Society of Pediatric Nephrology, Boston, MA (Lecture)
2000	Jansen PTH/PTHrP receptor in bone development.
	Molecular Medicine Seminar Series, April 17, University of Connecticut Health
	Center, CT (Visiting Professor)
2002	The role of HIF-1alpha and VHL in endochondral bone development.
	American Society for Matrix Biology, November 6-9, Houston, TX (Lecture)
2002	Hypoxia and HIF-1alpha in endochondral bone development.
	Endocrine Grand Rounds, Yale University, New Haven, CT (Visiting Professor)
2003	Molecular mechanisms of endochondral bone development.
	HSS-Cornell University, New York, NY (Visiting Professor)
2003	Hypoxia and HIF-1alpha in endochondral bone development.
	Amgen, Thousand Oaks, CA (Visiting Scientist)
2004	Hypoxia and HIF-1alpha in chondrogenesis

	Keystone Symposium on "Biology of Hypoxia: The role of oxygen sensing in development, normal function and disease (D3)", March 25-30, Steamboat Spring Colorado (Lecturer) Meeting Review in Genes and Dev., 2004, 18:2183-2194.
2005	Hypoxia and HIF-1alpha in endochondral bone development. Departmental Research Seminar Series, Department of Pathology, UAB,
	Birmingham, AL (Visiting Professor)
2005	Role of hypoxia and HIF-1alpha in chondrogenesis.
	Orthopaedic Research Seminar Series, March 23, Washington University, St. Louis, MO (Visiting Professor)
2005	Hypoxia and HIF-1alpha in endochondral bone development. Endocrine Grand Rounds, NIH, Bethesda, MD (Visiting Professor)
2005	Molecular mechanisms of endochondral bone development.
	Cedars Sinai UCLA, Los Angeles, CA (Visiting Professor)
2005	Hypoxia and HIF-1alpha in chondrogenesis
	UC Davis, Sacramento, CA (Visiting Professor)
2005	Hypoxia and HIF-1alpha in chondrogenesis
	NYAS Conference Skeletal Development and Remodeling, May 18-21, New York,
	NY (Seminar)
2006	PTH and the PTH/PTHrP receptor in bone stromal cells.
	UC Davis, Sacramento, CA (Visiting Professor)
2006	Molecular mechanisms of endochondral bone development: a transgenic approach.
	UC Davis, Sacramento, CA (Visiting Professor)
2006	Hypoxia and HIF-1alpha in chondrogenesis.
	Mesenchymal Stem Cell Biology: a Symposium, March 17, UCSF, San Francisco,
	CA (Seminar)
2006	Hypoxia and HIF-1alpha in chondrogenesis.
	Workshop on Growth Plate, June 11-15, Portland, OR (Seminar)
2007	Growth plate development and hypoxia.
	Meet the Professors, ASBMR, September 16-19, Honolulu (Seminar)
2007	Hypoxia and HIF-1alpha in chondrogenesis.
	Orthopaedic Grand Rounds, Thomas Jefferson University, Philadelphia, PA
	(Visiting Professor)
2008	The fetal growth place: a developmental model of cellular adaptation to hypoxia.
	Children Memorial Hospital, Northwestern University, June 20, Chicago, IL
	(Visiting Professor)
2008	The fetal growth plate: a developmental model of cellular adaptation to hypoxia.
	Department of Biochemistry, NJMC, Newark, NJ (Visiting Professor)
2009	Analysis of bone marrow stroma: lessons from mutant.
	Endocrine Grand Rounds, Yale University, New Haven, CT (Visiting Professor)
2009	Hypoxia and HIFs in chondrogenesis.
	FASEB Meeting, April 18-22, New Orleans, LA (Lecture)
2009	Hypoxia-dependent collagen modifications in limb bud development.
	NYAS Conference Skeletal Biology and Medicine, April 29-May 2, New York, NY
	(Seminar)
2010	Dual action of pVHL in limb bud mesenchyme.
	Keystone Symposia-hypoxia: Molecular Mechanisms of Oxygen Sensing and
	Response Pathways, January 19-24, Keystone, CO (Invited Short Talk)
2010	Oxygen sensing in cartilage and bone development.
	ASBMR, October 15-19, Toronto, Canada (Lecture)
2010	Oxygen sensing in cartilage and bone development.
	Rolanette and Berdon Lawrence Bone Disease Program of Texas Seminar Series.
	December 3, Houston, TX (Visiting Professor)
2012	HIFs and VHL in cartilage, bone and hematopoiesis.

	Louis V. Avioli Seminar Series, April 20, St. Louis, MO (Visiting Professor)
2012	Fibrocytes and marrow fibrosis.
	Meet-the-Professors, ASBMR, October 12-15, Minneapolis, MN (Invited Seminar)
2012	HIFs and VHL in cartilage, bone and hematopoiesis.
	October 12, Department of Medicine, UMI School of Medicine, Ann Arbor, MI
	(Visiting Professor)
2012	VHL in organogenesis.
-	4 th International Research Conference on Multiple Hereditary Exostoses. The
	Children's Hospital of Philadelphia Research Institute, November 1-4.
	Philadelphia, PA (Seminar)
2012	HIFs and VHL in cartilage, bone and hematopoiesis.
	November 15, Department of Chemical Engineering, USC, Columbia, SC (Visiting
	Professor)
2012	HIFs and VHL in cartilage, bone and hematopoiesis.
	December 6, Institute for Reproductive Health and Regenerative Medicine,
	Department of Pathology, KUMC, Kansas City, KS (Visiting Professor)
2013	Hypoxia signaling pathways in cartilage, bone and hematopoiesis: an update.
	Endocrine Seminar Series, November 1, University of Michigan, Ann Arbor, MI
	(Seminar)
2013	Hypoxia signaling pathways in cartilage and development and homeostasis.
	December 6, Lerner Institute, Cleveland Clinic, Cleveland, OH (Seminar)
2015	HIFs and VHL in cartilage and bone development: a role for angiogenesis.
	April 20, University of Delaware, Newark, DE (Seminar)
2017	Choke to start the engine: how impairment of mitochondrial function can improve
	survival of hypoxic cells in vivo. April 21, Medical School-Yale University, New
	Heaven, CT (Lecture)
2018	Enhancing mitochondrial respiration causes severe intracellular hypoxia and death
	of hypoxic cells. February 14, National Cancer Institute, NIH, Bethesda, MD
	(Lecture)
	Suppressing mitochondrial respiration is critical for hypoxia tolerance in the fetal
2018	growth plate. July 13, University of Little Rock, Little Rock, AR (Lecture)
0.001	Hypoxia and mitochondria in skeletal development. March 3 rd , University of
2021	Colorado, Denver, CO (Mack Clayton Lecture)
	Hypoxia and mitochondria in skeletal development. March 25 th , UCSF, San
2021	Francisco, CA (Lecture)
2022	How hypoxia and mitochondria shape up the skeleton. October 17 th , Northeastern
2022	University, Boston, MA (Lecture)
2024	Nitochondria, HIFs and the Control of Bone Mass. Annual Scientific Retreat
2024	Lawrence Family Bone Disease Program of Texas, Houston, TA (Keynote Lecture)
2024	Uxygen on the brink: now nypoxia and mitochondria shape the skeleton. UCIrvine,
	Irvine, CA (Invited Lecture)

International

1991	Cloning of the human PTH/PTHrP receptor. Istituto di Endocrinologia, Universita
	di Pisa, Italy
1994	The human PTH/PTH receptor: from cloning to human diseases. International
	Symposium on "Endocrinology under 35" Rome, Italy
1997	The PTH/PTHrP receptor in Jansen Metaphyseal Chondrodysplasia. Deutsche
	Gesellschaft Für Endokrinologie, May 6-8, Lübeck, Germany

1998	Constitutively active PTH/PTHrP receptors: in vivo and in vitro studies. Séminar des Chaire de Communications Cellulaires (Pr. J-P. Changeux) et de Médecine Expérimentale (Pr. P. Corvol) du Collége de France, February 26.27, Paris, France
1999	The PTH/PTHrP receptor in endochondral bone development. Workshop on
2000	Jansen PTH/PTHrP receptor in bone development. Workshop on Osteobiology, Wurzburg, Germany
2003	The PTH/PTHrP receptor in bone growth and remodeling. 30 th European Symposium on Calcified Tissues (ECTS) May 8-12 Rome Italy
2003	Hypoxia and HIF-1alpha in endochondral bone development. 1st Joint Meeting IBMS/IBMS_June 3-7_Osaka_Japan
2005	Hypoxia and HIF-1alpha in chondrogenesis. Gordon Conference on Cartilage, June 5-8 Il Ciocco Italy
2005	Hypoxia and HIF-1alpha in chondrogenesis. Gordon Conference on Bone and Teeth, July 10-15, The University of New England, Maine, USA
2006	Hypoxia and HIF-1alpha in chondrogenesis. 20 Th International Meeting of Biochemistry and Molecular Biology, June 19-23, Kyoto, Japan
2006	Hypoxia and HIF-1alpha in chondrogenesis. Research Symposium Developmental Biology in Orthopaedics. October 26-28. Toronto, Canada
2008	HIF-1alpha and chondrocytes: a tale of paradoxes. Shriners Hospital for Children- McGill University May 28 Montreal Canada
2009	PTH and bone marrow stroma. The 3 rd meeting of Bone and Cartilage Frontier, Tokyo Japan
2010	PTH and stem cells Parathyroids 2010 February 11-13 Pisa Italy
2011	HIFs and VHL in cartilage: a tale of paradoxes. Gordon Conference on Cartilage Piology and Pathology March 6 11, Venture, CA, USA
2011	Hypoxia signaling pathway in cartilage, bone and hematopoiesis. 3rd Joint Meeting IBMS/ECTS Athens, May 7-11, Greece
2012	Hypoxia signaling pathway in cartilage, bone and hematopoiesis Hypoxia 2012, April 4-5, Nantes, France
2013	Hypoxia signaling pathway in organogenesis. Dealing with Hypoxia: Regulatory Aspects in Cells, Tissues and Organisms, June 8-12, Oulu, Finland
2013	HIF-1alpha is essential for the development of the nucleus pulposus. Second International Research Meeting, "New Horizons in Intervertebral Disc Research," November 6-8, Philadelphia, PA, USA
2015	Fibrosis and HIF-1alpha-dependent tumors of the soft tissue upon loss of VHL in mesenchymal progenitors. The Tumor Microenvironment: 14 th International Workshop, August 27-29, Vancouver, British Columbia, Canada
2016	HIFs and the osteogenesis-angiogenesis coupling. Advances in Mineral Metabolism: International Workshop, March 27-31, Snowmass, CO, USA
2016	Hypoxia signaling pathways in bone. Meet-the-Professor, Advances in Mineral Metabolism: International Workshop, March 27-31, Snowmass, CO, USA
2016	HIFs and hypoxia in joint health and disease. Tackling Joint Disease by Understanding Crosstalk between Cartilage and Bone Research Symposium, April 28-30 Rosemont II, USA
2017	Impairment of mitochondrial respiration promotes survival of hypoxic cells. The Tumor Microenvironment: 15 th International Workshop, April 26-29, Miami Beach, FL, USA
2018	Choke to start the engine: how the impairment of mitochondrial function enables survival of hypoxic chondrocytes in vivo. Gordon Conference on Bone and Teeth, January 28-February 2. Galveston, TX, USA
2018	

2018	Growth plate development and closure from embryos to adult. AAOS/ORS The physis: Fundamental knowledge to a fantastic future through research. February 7- 9. Resempting ULSA
	Impairment of mitochondrial respiration promotes survival of HIF-1alpha deficient
2018	chondrocytes in vivo. Therapeutic targeting of hypoxia-sensitive
	pathways, Keystone Symposia, April 10-14, Oxford, UK
2010	The hypoxia signaling pathway in skeletal development. First annual fibrodysplasia
2019	ossificans progressive (FOP) and traumatic heterotopic ossification (HO)
2010	Symposium. September 21, University of Michigan, Ann Arbor, Mi Mitochondria and HIEs in skalatal davalonment. Cordon Research Conference on
2019	Cartilage March 17-22 Galveston TX USA
	Suppressing mitochondrial respiration is critical for hypoxia tolerance in the fetal
2021	growth plate. The Tumor Microenvironment: 16 th International Workshop, June 13-
	15, Miami Beach, FL, USA
2021	Hypoxia and Mitochondria in skeletal development. ECTS, May 6-8, Digital
	Congress
	Choke to start the engine: how the impairment of mitochondrial function enables
2022	survival of hypoxic chondrocytes. HYPOXYGEN Webinar "Effects of Hypoxia on
2022	the biology of tumors and normal cells", October 20
2022	Meet-the-Professor, ASBINK, September 8-12, Austin, 1A, USA
2025	November 20.22 Brugge Belgium
2023	How Hypoxia and Mitochondria Shane up the Skeleton Keystone Hypoxia: From
2025	Basic Mechanisms to Emerging Theranies May 28-31 Killarney KY Ireland
2024	How Hypoxia and Mitochondria Shape up the Skeleton. Gordon Research
_ • _ •	Conference on Collagen, July 9-24, New London, NH, USA.
	Mitochondria and the Control of Skeletal Development. Gordon Research
	Conference on Bone and Teeth, January 28-February 2, Galveston, TX, USA

Patents

Patent: 5,840,853 Date: Nov 24, 1998. Parathyroid Hormone receptor and DNA encoding the same receptor. I was member of the team who cloned the rat and opossum PTH/PTHrP receptor, and I then cloned the human homolog.

Bibliography

Completed Publications in Scientific Journals: Peer-Reviewed

- 1. Pacini F, Elisei R, Anelli S, Gasperini L, Schipani E (*I contributed to the analysis of the data*), Pinchera A. Circulating neuron-specific enolase in medullary thyroid cancer. The International Journal of Biological Markers 1986;1:85-88.
- Vitti P, Chiovato L, Lopez G, Lombardi A, Santini F, Mammoli C, Bassi P, Gryczynska M, Schipani E (*I contributed to the analysis of the data*) Tosti-Balducci M, Fenzi GF, Pinchera A. Measurement of TSAb directly in serum using FRTL-5 cells. J Endocrinol Invest 1988; 11:313-317.

- 3. Marcocci C, Pacini F, Elisei R, **Schipani E** (*I contributed to the collection and analysis of the data*), Ceccarelli C, Miccoli P, Arganin M, Pinchera A. Clinical and biological behavior of bone metastases from differentiated thyroid carcinoma. **Surgery 1989**; 106:960-966.
- 4. Jüppner H, Abou-Samra AB, Uneno S, **Schipani E** (*I performed some of the key experiments and I contributed to the analysis of the data*), Keutmann HT, Potts JT Jr, Segre GV. Properties of amino-terminal parathyroid hormone-related peptide modified at position 11-13. **Peptides 1990**; 11:1139-1142.
- Jüppner H, Abou-Samra AB, Freeman M, Kong XF, Schipani E (*I performed numerous of the critical experiments that led to the cloning of the opossun PTH/PTHrP receptor cDNA*), Richards J, Kolakowski LF, Hock F, Potts JT Jr, Kronenberg HM, Segre GV. A G protein-linked receptor for parathyroid hormone and parathyroid hormone-related peptide. Science 1991; 254:1024-1026.
- 6. Abou-Samra AB, Jüppner H, Force T, Freeman M, Kong XF, **Schipani E** (*I performed some of the critical experiments that led to the cloning of the rat PTH/PTHrP receptor cDNA*), Urena P, Richards J, Bonventre JV, Potts JT Jr, Kronenberg HM, Segre GV. Expression cloning of a common receptor for parathyroid hormone and parathyroid hormone-related peptide from rat osteoblast-like cells: A single receptor stimulates intracellular accumulation of both cAMP and inositol trisphosphates and increases intracellular free calcium. **Proc Natl Acad Sci USA 1992**; 89:2732-2736.
- 7. Hustmyer FG, **Schipani E** (*I contributed to the analysis of the data*), Peacock M. Bsml polymorphism at the parathyroid hormone receptor locus (PTHR) in three populations. **Hum Mol Genet 1993**; 2:1330.
- Schipani E, Karga H, Karaplis AC, Potts JT Jr, Kronenberg HM, Segre GV, Abou-Samra AB, Jüppner H. Identical complementary deoxyribonucleic acids encode a human renal and bone parathyroid hormone (PTH)/PTH-related peptide receptor. Endocrinology 1993;132(5):2157-2165. *This paper was the first report of the cloning of the human PTH/PTHrP receptor.*
- 9. Fukayama S, **Schipani E** (*I contributed to the design of the study, to the analysis of the data and to the writing of the manuscript*), Jüppner H, Lanske B, Kronenberg HM, Abou-Samra AB, Bringhurst FR. Role of protein kinase-A in homologous down-regulation of parathyroid hormone (PTH)/PTH-related peptide receptor messenger ribonucleic acid in human osteoblast-like SaOS-2 cells. **Endocrinology 1994**; 134:1851-1858.
- 10. Jüppner H, Schipani E (I performed numerous critical experiments, and I contributed to the design of the study and to the analysis of the data), Bringhurst FR, McClure I,

Keutmann HT, Potts JT Jr., Kronenberg HM, Abou-Samra AB, Segre GV, Gardella TJ. The extracellular, amino-terminal region of the Parathyroid Hormone (PTH)/PTH-related peptide receptor determines the binding affinity for carboxyl-terminal fragments of PTH(1-34). Endocrinology 1994; 134:879-884.

- Kong XF, Schipani E (I cloned and characterized the human gene encoding the PTH/PTHrP receptor), Lanske B, Joun H, Karperien M, Defize LHK, Jüppner H, Potts JT Jr., Segre GV, Kronenberg HM, Abou-Samra A-B. The rat, mouse and human genes encoding the receptor for parathyroid hormone and parathyroid hormone-related peptide are highly homologous. Biochem Biophys Res Commun 1994;200:1290-1299.
- 12. Gelbert L, Schipani E (*I contributed to the analysis of the data and to the writing of the manuscript*), Jüppner H, Abou-Samra A-B, Segre GV, Naylor S, Drabkin H, White R, Heath H Ill. Chromosomal localization of the parathyroid hormone/parathyroid hormone-related protein receptor gene to human chromosome 3p21.2-p24.2. J Clin Endocrinol Metab 1994; 79:1046-1048.
- 13. Schipani E, Hustmyer FG, Bergwitz C, Jüppner H. Polymorphism in exon M7 of the PTHR gene. Hum Mol Gen 1994; 3:1210.
- 14. Schipani E, Weinstein LS, Bergwitz C, Iida-Klein A, Kong XF, Stuhrmann M, Kruse K, Whyte MP, Murray T, Schmidtke J, van Dop C, Brickman AS, Crawford JD, Potts JT Jr., Kronenberg HM, Abou-Samra AB, Segre GV, Jüppner H. Pseudohypoparathyroidism type lb is not caused by mutations in the coding exons of the human parathyroid hormone (PTH)/PTH-related peptide receptor gene. J Clin Endocrinol Metab 1995; 80:1611-1621. *The study demonstrated that, differently from what had been for long time hypothesized, Pseudohypoparathyroidism type Ib, a rare endocrine disorder of calcium and phosphate homeostasis, is not caused by mutations in the PTH/PTHrP receptor gene. This finding prompted Dr. Harald Jueppner and Dr. Schipani to start a wide genome search that eventually led to identification of the Gsalpha gene as the gene critically involved in the pathogenesis of Pseudohypoparathyroidism type 1b (see below).*
- 15. Schipani E, Kruse K, Jüppner H. A constitutively active mutant PTH/PTHrP receptor in Jansen type metaphyseal chondrodysplasia. Science 1995;268:98-100. This paper was the first report of mutant, constitutively active PTH/PTHrP receptor in Jansen's metaphyseal chondrodysplasia, a severe form of short-limbed dwarfism associated to hypercalcemia. This discovery defined the cause of a devastating human disease, and showed the PTH/PTHrP receptor is a crucial regulator of endochondral bone development. Jansen's metaphyseal chondrodysplasia has been one of the first examples in the literature of a human disease being caused by a constitutively active G-protein coupled receptor. The mutant PTH/PTHrP receptors identified in Jansen's metaphyseal chondrodysplasia have been valuable tools for in vitro structure-functions studies directed to investigate how this receptor couples to and activates G-proteins. More importantly, they have allowed the

generation of transgenic mice that, in combination with knockout models, have defined the critical role of the PTH/PTHrP receptor in endochondral bone development.

- 16. Orloff JJ, Kats Y, Urena P, **Schipani E** (*I contributed to the analysis of the data*), Vasavada R, Philbrick W, Behal A, Abou-Samra AB, Segre GV, Jüppner H. Further evidence for a novel receptor for amino-terminal parathyroid hormone-related protein on keratinocytes and squamous carcinoma cell lines. **Endocrinology 1995**; 136:3016-3023.
- 17. Gardella TJ, Luck MD, Jensen GS, **Schipani** *E* (*I performed some of the key experiments and contributed to the analysis of the data*), Potts JT, Jueppner H. Inverse agonism of amino-terminally truncated parathyroid hormone (PTH) and PTH-related peptide (PTHrP) analogs revealed with constitutively active mutant PTH/PTHrP receptors. **Endocrinology 1996**; 137:3936-3941.
- Jüppner H, Schipani E. Receptors for parathyroid hormone and parathyroid hormonerelated peptide: from molecular cloning to definition of diseases. Curr Opin Nephrol 1996; 5:300-306.
- Schipani E, Langman CB, Parfitt AM, Jensen GS, Kikuchi S, Kooh SW, Cole WG, Jueppner H. Constitutively activated receptors for parathyroid hormone and parathyroid hormone-related peptide in Jansen's metaphyseal chondrodysplasia. N Engl J Med 1996; 335:708-714.
- 20. Parfitt AM, Schipani E (*I contributed to the analysis of the data and to the writing of the manuscript*), Rao DS, Kupin W, Han ZH, Jueppner H. Hypercalcemia due to constitutive activity of the Parathyroid Hormone (PTH)/PTH-related peptide receptor: Comparison with primary hyperparathyroidism. J Clin Endocrinol Metab 1996; 81:3584-3588.
- 21. Schipani E, Jensen GS, Pincus J, Nissenson RA, Gardella TJ, Jüppner H. Constitutive activation of the cyclic adenosine 3', 5'-monophosphate signaling pathway by parathyroid hormone (PTH)/PTH-related peptide receptors mutated at the two loci for Jansen's metaphyseal chondrodysplasia. Mol Endocrinol 1997; 11:851-858.
- 22. Schipani E, Lanske B, Hunzelman J, Luz A, Kovacs CS, Lee K, Pirro A, Kronenberg HM, Jüppner H. Targeted expression of constitutively active receptors for parathyroid hormone and parathyroid hormone-related peptide delays endochondral bone formation and rescues mice that lack parathyroid hormone-related peptide. Proc Natl Acad Sci USA 1997; 94:13689-13694. This study reported the generation and characterization of the first transgenic mouse model in which a constitutively active PTH/PTHrP receptor was expressed in vivo. This model significantly contributed to the understanding of the role of PTHrP and its receptor in endochondral bone development.

- 23. Jüppner H, Schipani E (*I contributed to the design of the experiments, and I performed some of them*), Bastepe M, Cole DEC, Lawson ML, Mannstadt M, Hendy GN, Plotkin H, Koshiyama H, Koh T, Crawford JD, Olsen BR, Vikkula M. The gene responsible for pseudohypoparathyroidism type Ib is paternally imprinted and maps in four unrelated kindreds to chromosome 20q13.3. Proc Natl Acad Sci USA 1998; 95:11798-803. The study was the first report of the genetic locus for Pseudohypoparathyroidism Ib a rare disorder of calcium and phosphate homeostasis. It was also the first demonstration that this disease is caused by a gene genetically imprinted. Future studies, which originated from this novel observation, demonstrated that deletions in the regulatory region of Gsalpha gene were the cause of Pseudohypoparathyroidism Ib.
- 24. Schipani E, Langman C, Hunzelman J, Le Merrer M, Loke KY, Dillon MJ, Silve C, Jüppner H. A novel Parathyroid Hormone (PTH)/PTH-related peptide receptor mutation in Jansen's metaphyseal chondrodysplasia. J Clin Endocrinol Metab 1999; 84:3052-3057.
- 25. Karp SJ, **Schipani E** (*I contributed to the design of the study and to the analysis of the data. I also performed some of the experiments described in the paper, and I generated and characterized one of the transgenic lines used in the study*), St-Jacques B, Hunzelman J, Kronenberg H, McMahon AP. Indian Hedgehog coordinates endochondral bone development via Parathyroid Hormone related-Protein-dependent and -independent pathways. **Development 2000**; 27:543-548.
- 26. Calvi ML, Sims NA, Hunzelman JL, Knight MC, Giovannetti A, Saxton JM, Kronenberg HM, Baron H, Schipani E. Activated parathyroid hormone/parathyroid hormone-related protein receptor in osteoblastic cells differentially affects cortical and trabecular bone. J Clin Invest 2001; 107: 277-286 (*Commentary on the paper in the same issue*). This paper reports the generation and characterization of a mouse transgenic line expressing a constitutively active PTH/PTHrP receptor in cells of the osteoblast lineage. The analysis of these transgenic mice has demonstrated that the PTH/PTHrP receptor is a crucial mediator of both bone forming and bone resorbing actions of PTH. Moreover, it has proven that the PTH/PTHrP receptor activity has different net effects in different bone compartments. These findings have underlined the complexity and heterogeneity of the osteoblast population and of their regulatory microenvironment.
- 27. Calvi ML, Schipani E. The PTH/PTHrP receptor in Jansen's metaphyseal chondrodysplasia. J Endocrinol Invest 2000; 23:545-554. (Invited)
- Long F, Schipani E. (I contributed to the design of the study and to the analysis of the data. I also performed some of the critical experiments reported in the paper), Asahara H, Kronenberg H, Montminy M. The CREB family of activators is required for endochondral bone development. Development 2001; 128:541-550.

- 29. Carter PH, Petroni B, Gensure R, **Schipani E** (*I contributed to the design of the study and performed some of the key experiments*), Potts JT, Jr, Gardella TJ. Selective and non-selective inverse agonists for constitutively active Type-1 Parathyroid Hormone Receptors: Evidence for altered receptor conformations. **Endocrinology 2001**; 142:1534-1545.
- Chung UI, Schipani E (I contributed to the design of the experiments; I also generated and characterized one of the mouse transgenic lines used in the study), McMahon AP, Kronenberg H. Indian Hedgehog couples chondrogenesis to osteogenesis in endochondral bone development. J Clin Invest 2001; 107:295-304. (Commentary on the paper in the same issue.)
- 31. Oliveria LM, Seminara B, Beranova M, Hayes FJ, Valkenburgh SB, **Schipani E.** (*I trained Dr. Oliveria in the use of TGGE (temperature gradient gel electrophoreses), which is a screening method to detect genomic mutations. I also contributed to the analysis of the data*), Costa EMF, Latronico AC, Crowley WF, Vallejo M. The importance of autosomal genes in Kallmann syndrome: genotype-phenotype correlations and neuroendocrine characteristics. J Clin Endocrinol Metab 2001; 86:1532-1538.
- 32. Beranova M, Oliveria LM, Bedecarrats GY, Schipani E (I trained Dr. Beranova in the use of TGGE (temperature gradient gel electrophoreses), which is a screening method to detect genomic mutations. I also contributed to the analysis of the data), Vallejo M, Ammini AC, Quintos JB, Hall JE, Martin KA, Hayes FJ, Pitteloud N, Kaiser UB, Crowley WF, Seminara SB. Prevalence, phenotypic spectrum, and modes of inheritance of gonadotropin-releasing hormone receptor mutations in idiopathic hypogonadotropic hypogonadism. J Clin Endocrinol Metab 2001; 86:1580-1588.
- 33. Imanishi Y, Hosokawa Y, Yoshimoto K, Schipani E (*I performed some of the experiments*), Mallya S, Papanikolau A, Kifor O, Tokura T, Sablosky M, Ledgard F, Gronowicz G, Wang TC, Schmidt EV, Hall C, Brown EM, Bronson R, Arnold A. Primary hyperparathyroidism caused by parathyroid-targeted overexpression of cyclin D1 in transgenic mice. J Clin Invest 2001; 107:1093-1102.
- 34. Gori F, **Schipani E** (*I contributed to the design of the experiments and to the analysis of the data*), Demay MB. Fibromodulin is expressed by both chondrocytes and osteoblasts during fetal bone development. **J Cell Biochem 2001**; 82:46-57.
- 35. Soegiarto DW, Kiachopoulos S, **Schipani E** (*I contributed to the analysis of the data and I generated and characterized one of the mouse transgenic lines used in the study*), Jüppner H, Erben R and Lanske B. Partial rescue of PTH/PTHrP receptor knockout mice by targeted expression of the Jansen transgene. **Endocrinology 2001**; 142:5303-5310.

- 36. Schipani E, Ryan HE, Didrickson S, Kobayashi T, Knight M and Johnson RS. Hypoxia in cartilage: HIF-1a is essential for chondrocyte growth arrest and survival. Genes Dev 2001; 15: 2865-2876. (Discussion on this paper in a "Research Roundup" in Journal of Cell Biology, 2001, 155, 693). This study was the first demonstration that fetal cartilage has an inner hypoxic region, and the transcription factor HIF-1 is essential for endochondral bone development.
- Kobayashi T, Chung UI, Schipani E (*I contributed to the analysis of the data; I generated and characterized one of the mouse transgenic line used in the study*), Starbuck M, Karsenty G, Katagiri T, Goad DL, Lanske B, and Kronenberg HM. PTHrP and Indian hedgehog control differentiation of growth plate chondrocytes at multiple steps. Development 2002; 129:2977-2986.
- 38. Schipani E. Conditional gene inactivation using cre recombinase. Calcified Tissue International 2002; 71:100-102. (Invited)
- 39. Pfander D, Cramer T, Schipani E (*I contributed to the analysis of the data and to writing the paper*), Johnson RS. HIF-1α controls extracellular matrix synthesis by epiphyseal chondrocytes. Journal of Cell Science 2003; 116 1819-1826.
- 40. Chiusaroli R, Maier A, Knight MC, Byrne M, Calvi L, Baron R, Krane SM, Schipani E. Collagenase Cleavage of Type I Collagen is essential for both Basal and Parathyroid Hormone (PTH)/PTH-Related Peptide Receptor- Induced osteoclast activation and has differential effects on discrete bone compartments. Endocrinology 2003; 144: 4106-4016.
- 41. Calvi LM, Adams GB, Weibrecht K, Weber JM, Olson DP, Knight MC, Martin RP, Schipani E (I generated and characterized the mouse transgenic line used in the study), Divieti P, Bringhurst FR, Milner LA, Kronenberg HM, Scadden DT. Osteoblastic cells regulate the haematopoietic stem cell niche. Nature 2003; 425: 841-846. This paper was the first demonstration the both PTH treatment and activation of the PTH/PTHrP receptor in osteoblasts leads to a significant expansion of the pool of hematopoietic stem cells both in vitro and in vivo. Furthermore, it provides evidence that the osteoblasts are critical component of the "niche" for hematopoietic stem cells. She generated and characterized the mouse transgenic line used in this study; this mouse line was of critical importance for the design and success of the study.
- 42. Schipani E, Provot S. PTHrP, PTH and the PTH/PTHrP receptor in endochondral bone development. Birth Defects Research 2003; 69:352-362. (Invited)
- 43. Strowski MZ, Cramer T, Schafer G, Juttner S, Walduck A, Schipani E (I performed the in

situ hybridization analysis), Kemmer W, Wessler S, Wunder C, Weber M, Meyer TF, Wiedenmann B, Jons T, Naumann M, Hocker M. Helicobacter Pylori stimulates host Vascular Endothelial Growth Factor-A (vegf-A) gene expression via MEK/ERK-dependent activation of Sp1 and Sp3. FASEB J 2004; 18:218-220.

- 44. Calvi LM, Shin HI, Knight MC, Weber JM, Young M.F, Giovannetti A, Schipani E. Constitutively active PTH/PTHrP receptor in odontoblast and ameloblast function and maturation. Mechanisms of Development 2004; 121:397-408.
- 45. Zelzer E, Mamluk R, Ferrara N, Johnson RS, **Schipani E** (*I contributed to design of the study, to the analysis of the data, and to writing the mansucript; I also performed some of the critical experiments shown in the paper*), and Olsen BR. VEGFA is necessary for chondrocyte survival during bone development. **Development 2004**; 131:2161-2171.
- 46. Pfander D, Kobayashi T, Knight MC, Zelzer E, Chan DA, Olsen BR, Giaccia AJ, Johnson RS, Haase V, Schipani E. Deletion of Vhlh in chondrocytes reduces cell proliferation and increases matrix deposition during growth plate development. Development 2004; 131:2497-2508. This study was the first in vivo demonstration that loss of the tumor suppressor gene VHL (Von Hippel Lindau) paradoxically inhibits chondrocyte proliferation during fetal development, and causes severe dwarfism.
- 47. Cramer T, Schipani E (*I contributed to the design of the experiments, analysis of the data and writing of the paper*), Johnson RS, Swodoba B, Pfander D. Expression of VEGF isoforms by epiphyseal chondrocytes during low-oxygen tension is HIF-1α dependent. Osteoarthritis Cartilage 2004; 12:433-439.
- 48. Larsson T, Marsell R, Schipani E (I contributed to the design of the study and to the analysis of the data; I also performed some of the experiments shown in the paper), Ohlsson C, Ljunggren O, Tenenhouse HS, Jüppner H and Jonsson K. Transgenic mice expressing Fibroblast Growth Factor 23 under the control of the α1(I) collagen promoter exhibit growth retardation, osteomalacia and disturbed phosphate homeostasis. Endocrinology 2004; 7:3087-3094.
- 49. Fröhlich LF, Gensure RC, Schipani E (*I contributed to the analysis of the data and to writing the paper*), Jüppner H, Bastepe M. Haplotype frequencies and linkage disequilibrium analysis of four frequent polymorphisms at the PTH/PTH-related peptide receptor gene locus. Mol Cell Probes 2004; 18: 53-357.
- 50. Kuznetsov SA, Riminucci M, Ziran N, Tsutsui TW, Corsi A, Calvi L, Kronenberg HM, **Schipani E** (*I contributed to the design of the study, to the analysis of the data and to writing the paper; I provided the mouse transgenic line used in the study and performed key*

experiments), Robey PG, Bianco P. The interplay of osteogenesis and hematopoiesis: expression of a constitutively active PTH/PTHrP receptor in osteogenic cells perturbs the establishment of hematopoiesis in bone and of skeletal stem cells in the bone marrow. J Cell Biol 2004; 167:1113-1122.

- 51. Kobayashi T, Soegiarto DW, Yang Y, Lanske B, **Schipani E** (*I contributed to the analysis of the data and generated one of the mouse transgenic models used in the study*), McMahon AP, Kronenberg HM. Indian Hedgehog stimulates periarticular chondrocyte differentiation to regulate growth plate length independently of PTHrP. J Clin Invest 2005 115:1734-1742
- 52. Schipani E. Hypoxia and HIF-1a in chondrogenesis. Seminars in Cell Biology and Development 2005; 16:539-546. (Invited)
- 53. Provot S, Schipani E. Molecular mechanisms of endochondral bone development. Biochem Biophys Res Commun 2005; 328:658-65. (Invited)
- 54. Carter PH, Schipani E. The Roles of Parathyroid Hormone and Calcitonin in Bone Remodeling: Prospects for Novel Therapeutics. Endocrine, Metabolic and Immune Disorders- Drug Targets 2006; 6: 59-76. (Invited)
- 55. Schipani E. Hypoxia HIF-1a and VHL in chondrogenesis. Annals of New York Academy of Sciences 2006; 1068: 66-73. (Invited)
- 56. Zambetti GP, Horwitz E, **Schipani E**. Skeletons in the p53 tumor suppressor closet: genetic evidence that p53 blocks bone differentiation and development. **J Cell Biol 2006**; 172: 795-797. (Invited)
- 57. Marsell R, Jonsson KB, Cho TJ, Einhorn TA, Ohlsson C, **Schipani E**. Fracture healing is not impaired in mice expressing a constitutively active PTH/PTHrP receptor in osteoblasts. Acta Orthopaedica 2007; 78:39-45
- 58. Provot S, Zinyk D, Gunes Y, Khatri R, Le Q, Longaker MT, Giaccia AJ, Schipani E. Hifla regulates differentiation of limb bud mesenchyme and joint development. J Cell Biol 2007; 177:451-464. (Cover Figure and Commentary in the same issue). This study was the first in vivo demonstration that HIF-1 is not only a survival factor but also a differentiation factor for the developing cartilage, as its loss in limb bud mesenchyme considerably delays both the timely transition of mesenchymal cells into chondrocytes and joint development.
- 59. Ono N, Nakashima K, **Schipani E** (*I contributed to the analysis of the data and generated one of the mouse transgenic models used in the study*), Hayata T, Ezura Y Soma K,

Kronenberg HM, Noda M. Constitutively active parathyroid homone signaling in cells in osteoblastic lineage suppresses mechanical unloading-induced bone resorption. **J Biol Chem 2007**; 282: 25509-25516.

- 60. Wang Y, wan C, Ximeng L, Bouxsein ML, Faugere MC, Guldberg RE, Johnson RS, Haase VH, Gerstenfield LC, Schipani E (*I designed the study together wit Dr. Clemens, I contributed to the analysis of the data and to writing the manuscript; I performed critical experiments*), Clemens TL. The hypoxia inducible factor pathway couples angiogenesis to osteogenesis during skeletal development. J Clin Invest 2007, 117: 1616-1626 (*Cover Figure and Commentary in the same issue*). This study was the first demonstration in vivo that activation of the hypoxia signaling pathway in terminally differentiated osteoblasts dramatically increases bone mass and number of bone marrow blood vessels. The paper strongly supported the notion that the osteogenesis-angiogenesis coupling is likely to be a crucial event in bone development and homeostasis.
- 61. Provot S, Schipani E. Fetal growth plate: a developmental model of cellular adaptation to hypoxia. Annals of New York Academy of Sciences 2007; 1117:26-39. (Invited)
- 62. Neri P, Tassone P, Shammas M, Yasui H, **Schipani E** (*I performed the in situ hybridization analysis shown in the paper*), Batchu RB, Blotta S, Prabhala R, Catley L, Hamasaki M, Hideshima T, Chauhan D, Jacob GS, Picker D, Venuta S, Anderson KC, Munshi NC. Biological pathways and in vivo antitumor activity induced by Atiprimod in myeloma. Leukemia 2007, 21: 2519-2526.
- 63. Kobayashi T, Lu J, Cobb BS, Rodda SJ, McMahon AP, **Schipani E** (*I contributed to the analysis of the data; I generated and characterized one of the mouse transgenic line used in the study*), Merkenschlager M, Kronenberg HM. Dicer-dependent pathways regulate chondrocyte proliferation and differentiation. **PNAS 2008**, 105: 1949-1954.
- 64. Ono N, Nakashima K, Rittling SR, **Schipani E** (*I contributed to the analysis of the data and to writing the paper; I generated and characterized one of the mouse transgenic line used in the study*), Hayata T, Ezura Y, Soma K, Denhardt DT, Kronenberg HM, Noda M. Osteopontin negatively regulates Parathyroid Hormone Receptor Signaling in Osteoblasts. **JBC 2008**, 283: 19400-19409.
- 65. Larsson J, Ohishi M, Garrison B, Aspling M, Janzen V, Adams GB, Curto M, McClatchey AI, **Schipani E** (*I contributed to the design of the study, the analysis of the data and the writing of the manuscript; I performed some of the experiments shown in the study),* Scadden DT. Cell contact growth inhibitor Nf2/merlin regulates stem cell behavior by altering microenvironmental architecture. **Cell Stem Cell 2008**, 3: 221-227.

- 66. Shimada M, Greer PA, McMahon AP, Bouxsein ML, Schipani E. In vivo targeted deletion of Capn4 in cells of the osteoblast lineage impairs osteoblast function and bone formation. JBC 2008, 283: 21002-21010.
- 67. O'Brien CA, Plotkin LI, Galli C, Goellner JJ, Gortazar AR, Robling A, Bouxsein M, Schipani E (*I contributed to the analysis of the data and the writing of the manuscript; I performed some of the experiments shown in the study*),, Turner CH, Jilka RL, Weinstein RS, Manolagas SC, Bellido T. Control of bone mass and remodeling by PTH receptor signaling in osteocytes. **PloSOne 2008**, 3: E2942.
- 68. Schipani E. A new window on MSC (Commentary). Blood 2008; 112: 217-218. (Invited)
- 69. Khatri R, Schipani E. About the importance of being desulfated (Commentary). Genes Dev 2008; 22(20):2750-2754. (Invited)
- 70. Ohishi M, Chiusaroli R, Thomas C, Asuncion F, Ominksy M, Kostenuik P, **Schipani E.** Osteoprotegerin cures bone marrow fibrosis and cortical porosity in a mouse model of constitutive activation of the PTH/PTHrP receptor. **Am J Pathol 2009**; 174:2160-2171
- 71. Maeda Y, Schipani E (I contributed to the design of the study, the analysis of the data and the writing of the manuscript; I performed some of the experiments shown in the study), Densmore MJ, Lanske B. Partial rescue of postnatal growth plate abnormalities in Ihh mutants by expression of a constitutively active PTH/PTHrP receptor. Bone 2009; 46: 472-478.
- 72. Schipani, E and Kronenberg, H.M. Adult mesenchymal stem cells (January 31, 2009), StemBook, The Stem Cell Research Community, StemBook, doi/10.3824/stembook.1.38.1. (Invited)
- Riddle RC, Khatri R, Schipani E (*I contributed to the writing of the manuscript*), Clemens TL. Role of hypoxia –inducible –factor 1alpha in angiogenic-osteogenic coupling. J Mol Med, 2009; 87: 583-590. (Invited)
- 74. Schipani E, Maes C, Carmeliet G, Semenza GL. Regulation of osteogenesis-angiogenesis coupling by HIFs and VEGF. J Bone Miner Res, 2009; 24: 1347-1353. (Invited)
- 75. Myllyharju J, Schipani E. Extracellular matrix genes and hypoxia-inducible targets. Cell and Tissue Research, 2010; 339:19-29. (Invited)
- 76. Ohishi M, Schipani E. Bone Marrow Mesenchymal Stem cells. J Cell Biochem 2010;

109:277-282. (Invited)

- 77. Risbud MV, Schipani E (*I contributed to the writing of the manuscript*), Shapiro IM. Hypoxic Regulation of Nucleus Pulposus Cell Survival. From Niche to Notch. Am J Pathol 2010; 176: 11577-11583.
- 78. Guo J, Liu M, Yang D, Bouxsein ML, Saito H, Galvin RJ, Kuhstoss SA, Thomas CC, Schipani E (I contributed to the design of the study and to the analysis of the data; I generated and characterized one the transgenic lines used in this study), Baron R, Bringhurst FR, Kronenberg HM. Suppression of Wnt signaling by Dkk1 attenuates PTHmediated stromal cell response and new bone formation. Cell Metab. 2010 Feb 3; 11(2):161-71.
- 79. Schipani E. Posttranslational modifications of collagens as targets of hypoxia and Hiflalpha in endochondral bone development Annals of New York Academy of Sciences 2010; 1192: 317-321. (Invited)
- Wan C, Shao J, Gilbert SR, Riddle RC, Long F, Johnson RS, Schipani E, Clemens TL. Role of Hif-1alpha in skeletal development. Annals of New York Academy of Sciences 2010; 1192: 322-326. (Invited)
- 81. Giaccia AJ and Schipani E. Role of carcinoma-associated fibroblasts and hypoxia in tumor progression. Curr Top Microbiol Immunol 2010; 810: 31-45. (Invited)
- Kashigawi A, Schipani E (I trained Dr. Kashigawi in the histology techniques, I contributed to the design of the study, to the analysis of the data and to writing of the manuscript), Fein MJ, Greer PA, Shimada M. Targeted deletion of Capn4 in cells of the chondrocyte lineage impairs chondrocyte proliferation and differentiation. Mol Cell Biol 2010; 30: 2799-2810.
- 83. Guo J, Liu M, Yang D, Bouxsein ML, Thomas CC, **Schipani E** (*I contributed to the design of the study and to the analysis of the data; I generated and characterized one the transgenic lines used in this study*), Bringhurst FR, Kronenberg HM. Phospholipase C signaling via the Parathyroid Hormone (PTH)/PTH-related peptide receptor is essential for normal bone response to PTH. **Endocrinology 2010**; 151: 3502-3513.
- 84. Araldi E and Schipani E. Hypoxia, Hifs and bone development. Bone, 2010; 47: 190-196. (Invited)

- 85. Araldi E and Schipani E. microRNA-140 and the silencing of osteoarthritis. Genes Dev 2010; 24: 1075-1080. (Invited)
- 86. Hanyu R, Hayata T, Nagao M, SaitaY, Hemmi H, Notomi T, Nakamoto T, Schipani E (*I contributed to the analysis of the data; I generated and characterized one the transgenic lines used in this study*), Kronenberg H, Kaneko K, Kurosawa H, Ezura Y, Noda M.Per-1 is a specific clock gene regulated by parathyroid (PTH) signaling in osteoblasts and isfunctional for transcriptional events induced by PTH. J Cell Biochem 2011; 122: 433-8.
- 87. Araldi E, Khatri R, Giaccia AJ, Simon MC, Schipani E. Lack of Hif-2alpha in limb bud mesenchyme causes only a modest and transient delay of endochondral bone development. Nature Medicine 2011;17:25-26. This paper was the first in vivo demonstration that differently from HIF-1, which has an essential and non-redundant role in endochondral bone development, HIF-2, another member in the family of HIF transcription factors, is dispensable for fetal cartilage. Notably, HIF-2 is thought to have an important role in the homeostasis of articular surface cartilage. This study thus supports the notion that fetal growth plate cartilage and adult articular surface cartilage are distinct biological entities.
- 88. Rankin EB, Giaccia AJ, Schipani E. A central role for hypoxic signaling in cartilage, bone and hematopoiesis. Curr Osteoporos Rep. 2011; 9:46-52.
- 89. Ono N, Nakashima K, **Schipani E** (*I contributed to the design of the study and to the analysis of the data; I generated and characterized one the transgenic lines used in this study*), Hayata T, Ezura Y, Soma K, Kronenberg HM, Noda M. Constitutively active PTH/PTHrP receptor specifically expressed in osteoblasts enhances bone formation induced by bone marrow ablation. **J Cell Physiol 2012** Feb;227(2):408-15.
- 90. Ohishi M, Ono W, Ono N, Khatri R, Marzia M, Baker EK, Root SH, Wilson TLS, Iwamoto Y, Kronenberg HM, Aguila HL, Purton LE, Schipani E. A novel population of cells expressing both hematopoietic and mesenchymal markers is present in the normal adult bone marrow and is augmented in a murine model of marrow fibrosis. Am J Pathol 2012;180:811-8 (*Highlight in AJP*). This paper reported for the first time the presence in the normal bone marrow of cells that express both hematopoietic cell surface antigens such as CD45 and CD11b, and classical mesenchymal markers such as collagen type I, and are thus phenotypically similar to fibrocytes. Fibrocytes are present in wounds, at sites of pathological fibroses, and in the reactive stroma of tumors. Notably, this novel bone marrow is increased by PTH treatment.
- 91. Rankin EB, Wu C, Khatri R, Wilson TLS, Rankin AB, Kuo CJ, **Schipani E.** (*I designed the study together with Dr. Giaccia; I trained Dr. Rankin and Dr. Wu in the analysis of bone phenotypes in vivo; I analyzed all the data together with Drs. Giaccia, Rankin and Wu, and together with them I wrote the manuscript; I generated and characterized one the conditional knockouts described in the paper that have been essential for the success of the success o*

study), Giaccia AJ. HIF signaling in osteoblasts regulates erythroid progenitors through EPO. Cell 2012; 149: 63-74. This study revealed an unexpected role for osteoblasts in the production of EPO and modulation of erythropoiesis. Furthermore, it demonstrated a molecular role for the HIF signaling pathway in osteoblasts that can be targeted to elevate both hematopoietic stem cells and erythroid progenitors in the local hematopoietic microenvironment, and to dramatically increase bone mass.

- 92. Maes C, Araldi E, Haigh K, Khatri R, Van Looeveren A, Giaccia AJ, Haigh JJ, Carmeliet G, Schipani E. VEGF-independent cell –autonomous functions of HIF-1alpha regulating oxygen consumption in fetal cartilage are critical for chondrocyte survival. JBMR 2012; 27:596-609. This paper demonstrated for the first time that VEGF, though it is both a survival factor for chondrocytes and a classical downstream target of HIF-1 transcriptional activity, has only a modest role, if any, in mediating the survival function of HIF-1 in cartilage. In particular, HIF-1 null chondrocytes are dramatically hypoxic in vivo. Since the degree of oxygenation of a cell is the net result of O_2 availability and O_2 consumption, we asked whether loss of HIF-1 in chondrocytes affects any of these two variables or both. We discovered that loss of VEGF in chondrocytes causes chondrocyte death, decreases number of blood vessels in the surrounding soft tissue, and increases degree of hypoxia of viable chondrocytes. VEGF overexpression in chondrocytes fully corrected inner cell death and increased hypoxia of VEGF null chondrocytes, most likely by augmenting number of blood vessels in the surrounding soft tissue. Based on these observations, we concluded that VEGF is a survival factor for chondrocytes mostly likely by increasing O₂ availability to these cells. Surprisingly, however, VEGF overexpression in chondrocytes only minimally corrected inner cell death and increased hypoxia of HIF-1 null growth plates, despite a dramatic increase of number of blood vessels in the surrounding soft tissue. Taken altogether, VEGF is unlikely to be the only mediator of the HIF-1 role as a survival factor; moreover, decreased O_2 availability is unlikely to be the only cause of the severe hypoxia of viable HIF-1 null chondrocytes. Thus, if O_2 availability is not the key element, O_2 consumption has to be the essence of the problem. It is possible that the main tool adopted by HIFs to keep in check hypoxia is indeed cell autonomous control of metabolism. It is now established that, at least in vitro, HIF-1 stimulates non-oxidative glycolysis and impairs oxidative phosphorylation mainly by inhibiting entry of pyruvate into the Krebs cycle .It is possible that HIF-1-dependent impairment of mitochondrial respiration and thus of O_2 consumption is critical to keep in check hypoxia and thus allow chondrocyte survival. We are currently testing this hypothesis.
- 93. Maes C, Carmeliet G, Schipani E. Hypoxia signaling pathway and angiogenesis in bone development, regeneration and disease. Nature Review Rheumatology 2012; 8:358-366.
- 94. Aro E, Khatri R, Gerard-O'Riley R, Myllyharju J and **Schipani E**. Hypoxia-inducible factor-1 (HIF-1) but not HIF-2 is essential for hypoxic induction of collagen prolyl 4-hydroxylases in mouse epiphyseal growth plate chondrocytes. **JBC 2012**; 287:37134-44.

This study investigated the role of hypoxia and HIFs in regulating expression of collagen prolyl 4-hydroxylases (C-P4Hs) in chondrocytes in vitro. Briefly, C-P4Hs are key enzymes in collagen synthesis as the resulting 4-hydroxyprolines are necessary for the stability of all collagen molecules. C-P4H I is the main form in most cells, but C-P4H II is the major form in chondrocytes. We utilized primary epiphyseal growth plate chondrocytes isolated from newborn mice with conditionally inactivated HIF-1 or HIF-2 genes, and we showed that C-P4H I and II mRNA levels were increased in hypoxic chondrocytes, the induction being HIF-1 but not HIF-2 dependent. Furthermore, the increases in the C-P4H mRNA levels were associated with increased amounts of the C-P4H proteins and C-P4H activity in hypoxia. Taken together, our findings indicate that the hypoxia-inducibility of the C-P4H isoenzymes is likely to ensure sufficient C-P4H activity for collagen synthesis occurring in chondrocytes in a hypoxic environment. These findings demonstrated for the first time that HIF-1 and not HIF-2 is a critical regulator of collagen synthesis in chondrocytes.

- 95. Hanyu R, Wehbi VL, Hayata T, Moriya S, Feinstein TN, Ezura Y, Nagao M, Saita Y, Hemmi H, Notomi T, Nakamoto T, Schipani E (*I contributed to the design of the study and to the analysis of the data; I generated and characterized one the transgenic lines used in this study*), Takeda S, Kaneko K, Kurosawa H, Karsenty G, Kronenberg HM, Vilardaga JP, Noda M. Anabolic action of parathyroid hormone regulated by the β2-adrenergic receptor. Proc Natl Acad Sci U S A 2012; 109:7433-8.
- 96. Simic P, Zainabadi K, Bell E, Sykes DB, Saez B, Lotinun S, Baron R, Scadden D, Schipani E (*I performed the histological analysis reported in the study*), Guarente L. SIRT1 regulates differentiation of mesenchymal stem cells by deacetylating β-catenin. EMBO Mol Med. 2013; 5:430-40.
- 97. Krause D, Fulzele K, Catic A, Chia-Chi S, Dombkwoski D, Hurley MP, Lezeau S, Attar E, Wu JY, Lin HY, Divieti-Pajevic P, Hasserjian, **Schipani E** (*I contributed to the design of the study, to the analysis of the data, and to writing the paper; I generated and characterized one of the transgenic lines used in this study*), Van Etten RA, Scadden DT. Differential regulation of myeloid leukemias by the bone marrow microenvironment. **Nature Medicine 2013;** 19:1513-7.
- 98. Schipani E, Wu C, Rankin BE and Giaccia A. Role of osteoblasts in regulating bone marrow angiogenesis during bone development and homeostasis. Front. Endocrinol 2013 Jul 10;4:85.
- 99. Schipani E, Mangiavini L and Merceron C. ATF4 and HIF-1 in bone: an intriguing relationship. JBMR 2013; 28:1866-9.
- 100. Mangiavini L, Araldi E, Khatri R, Wilson TLS, Gerard-O'Riley R, Rankin EB, Giaccia AJ and Schipani E. Loss of VHL in mesenchymal cells of the limb bud alters multiple steps of endochondral bone development. Dev Biol 2014; 393(1):124-36. This study investigated the role of VHL in mesenchymal progenitor cells of the limb bud and their descendants. Briefly,

loss of VHL in limb bud mesenchyme did not alter the timely differentiation of mesenchymal cells into chondrocytes. However, it caused structural collapse of the cartilaginous growth plate as a result of impaired proliferation, delayed terminal differentiation, and ectopic death of chondrocytes. This phenotype was associated to delayed replacement of cartilage by bone. Our findings demonstrate that VHL regulates bone morphogenesis as its loss considerably alters size, shape and overall development of the skeletal elements.

- 101. Merceron C, Mangiavini L, Wilson TLS, Robling A, Giaccia AJ, Shapiro I, Risbud M and Schipani E. Loss of HIF-1a in the notochord results in cell death and complete disappearance of the nucleus pulposus. PlosOne 2014; 9(10):e110768. The intervertebral disc (IVD) is one of the largest avascular organs in vertebrates. The nucleus pulposus (NP), a highly hydrated and proteoglycan-enriched tissue, forms the inner portion of the IVD. The NP is surrounded by a multi-lamellar fibrocartilaginous structure, the annulus fibrosus (AF). This structure is covered superior and inferior side by cartilaginous endplates (CEP). The NP is a unique tissue within the IVD as it results from the differentiation of notochordal cells, whereas, AF and CEP derive from the sclerotome. The hypoxia inducible factor-1a (HIF-1a) is expressed in NP cells but its function in NP development and homeostasis is largely unknown. We thus conditionally deleted HIF-1a in notochordal cells and investigated how loss of this transcription factor impacts NP formation and homeostasis at E15.5, birth, 1 and 4 months of age, respectively. Histological analysis, cell lineage studies, and TUNEL assay were performed. Morphologic changes of the mutant NP cells were identified as early as E15.5, followed, postnatally, by the progressive disappearance and replacement of the NP with a novel tissue that resembles fibrocartilage. Notably, lineage studies and TUNEL assay unequivocally proved that NP cells did not transdifferentiate into chondrocyte-like cells but they rather underwent massive cell death, and were completely replaced by a cell population belonging to a lineage distinct from the notochordal one. Finally, to evaluate the functional consequences of HIF-1a deletion in the NP, biomechanical testing of mutant IVD was performed. Loss of the NP in mutant mice significantly reduced the IVD biomechanical properties by decreasing its ability to absorb mechanical stress. These findings are similar to the changes usually observed during human IVD degeneration. Our study thus demonstrates that HIF-1a is essential for NP development and homeostasis, and it raises the intriguing possibility that this transcription factor could be involved in IVD degeneration in humans.
- 100. Johnson RW, Schipani E and Giaccia AJ. HIF targets in bone remodeling and metastatic disease. Pharmacology and Therapeutics 2015; S0163-7258(15)00046-7.
- 101. Wu Colleen, Rankin EB, LaGory EL, Andersen R, Rhodes SD, Wilson TLS, Mohammed KS, Castillo A, Guise TA, **Schipani E** (*I contributed to the design of the study, to the generation and analysis of the data, and to writing the paper*), Giaccia AJ. Selective inactivation of osteoblastic HIF prolyl hydroxylases regulates bone homeostasis through the regulation of OPG. **Genes Dev 2015**; 29: 791-802.
- 102. Aro E, Salo AM, Khatri R, Finnila M, Miinalainen I, Sormunen R, Pakkanen O, Holster T, Soininen R, Prein C, Clausen-Schaumann H, Aszodi A, Tuukannen J, Kivirikko KI, Schipani E* and Myllyharju J*. Severe extracellular matrix abnormalities and chondrodysplasia in mice lacking collagen prolyl-4-hydroxylase isoenzyme II in

combination with a reduced amount of isoenzyme I. (* co-senior authors) **JBC 2015**; 90:16964-16978.

- 103. Schipani E, Mangiavini L, Merceron C. HIF-1a and the developing growth plate: what we really now. BoneKey 2015; 30. doi: 10.1038/bonekey.2015.99. eCollection 2015 (Invited).
- 104. Mangiavini L, Merceron C, Araldi E, Khatri R, Gerard-O'Riley R Wilson TLS, Sandusky G. Abadie J, Lyons LM, Giaccia AJ and Schipani E. Fibrosis and HIF-1a-dependent tumors of the soft tissue upon loss of VHL in mesenchymal progenitors. AJP 2015; 185:3090-101. Loss of Vhl in mesenchymal progenitors of the limb bud caused severe fibrosis of the synovial joints and formation of aggressive masses with histologic features of mesenchymal tumors. Hif-1α and its downstream target connective tissue growth factor were necessary for the development of these tumors, which conversely still developed in the absence of Epas1, but at lower frequency. Human tumors of the soft tissue are a very complex and heterogeneous group of neoplasias. Our novel findings in genetically altered mice suggest that activation of the HIF signaling pathway could be an important pathogenetic event in the development and progression of at least a subset of these tumors.
- 105. Pineault KM, Swinehart IT, Garthus KN, Ho E, Yao Q, **Schipani E** (*1 performed the ISH analysis on histological sections*), Kozloff KM, Wellik DM. Hox11 genes regulate postnatal longitudinal bone growth and growth plate proliferation. **Biol Open. 2015**; 4(11):1538-48.
- 106. Sinha P, Aarnisalo P, Chubb R, Poulton IJ, Guo J, Nachtrab G, Kimura T, Swami S, Saeed H, Chen M, Weinstein LS, **Schipani E** (*I generated one of the mouse transgenic lines used in he paper; I edited the manuscript*), Sims NA, Kronenberg HM, Wu JY. Loss of Gsα in the Postnatal Skeleton Leads to Low Bone Mass and a Blunted Response to Anabolic Parathyroid Hormone Therapy. **JBC 2016**; 291:1631-42.
- 107. Agarwal S, Loder S, Brownley C, Cholok D, Mangiavini L, Li J, Breuler C, Sung HH, Li S, Ranganathan K, Peterson J, Tompkins R, Herndon D, Xiao W, Jumlongras D, Olsen BR, Davis TA, Mishina Y, Schipani E (*co-corresponding author), Levi B. Inhibition of Hiflα prevents both trauma-induced and genetic heterotopic ossification. Proc Natl Acad Sci U S A 2016;113:E338-47. Pathologic extraskeletal bone formation, or heterotopic ossification (HO), occurs following mechanical trauma, burns, orthopedic operations, and in patients with hyperactivating mutations of the type I bone morphogenetic protein receptor ACVR1 (Activin type 1 receptor). Extraskeletal bone forms through an endochondral process with a cartilage intermediary prompting the hypothesis that hypoxic signaling present during cartilage formation drives HO development and that In this study, we demonstrated that Hifla is highly expressed and active in three separate mouse models: trauma-induced, genetic, and a hybrid model of genetic and trauma-induced HO. Pharmacologic inhibition of Hif1a using PX-478 or rapamycin significantly decreased or inhibited extraskeletal bone formation. Importantly, de novo soft-tissue HO was eliminated or significantly diminished in treated mice. Lineage-tracing mice demonstrated that cells forming HO belong to the Prx lineage. Burn/tenotomy performed in lineage-specific Hif1a knockout mice resulted in

substantially decreased HO, and again lack of de novo soft-tissue HO. Genetic loss of Hif1a in mesenchymal cells marked by Prx-cre prevented the formation of the mesenchymal condensations. Pharmacologic inhibition of Hif1a had a similar effect on mesenchymal condensation development. Our findings indicate that Hif1a represents a promising target to prevent and treat pathologic extraskeletal bone.

- 108. Wang H, Lindborg C, Lounev V, Kim JH, McCarrick-Walmsley R, Xu M, Mangiavini L, Groppe JC, Shore EM, Schipani E (*I provided critical reagents, I contributed to the design of the study and to the interpretation of the data and I critically reviewed the manuscript*), Kaplan FS and Pignolo RJ. Cellular Hypoxia Promotes Heterotopic Ossification by Amplifying BMP Signaling. J Bone Miner Res. 2016; 31:1652-65.
- 109. Choi H, Merceron C, Mangiavini L, Seifert EL, Schipani E (I provided critical reagents and I critically reviewed the manuscript), Shapiro IM and Risbud MV. Hypoxia promotes noncanonical autophagy in nucleus pulposus cells independent of MTOR and HIF1A signaling. Autophagy 2016;12:1631-46.
- 110. Mirzamohammadi F, Papaioannou G, Inloes JB, Rankin EB, Xie H, Schipani E (*I provided critical reagents, I contributed to the design of the study and to the interpretation of the data and I critically reviewed the manuscript*), Orkin SH and Kobayashi T. Polycomb repressive complex 2 regulates skeletal growth by suppressing Wnt and TGF-β signalling. Nat Commun. 2016;7:12047.
- 111. Logsdon DP, Grimard M, Luo M, Shahda S, Jiang Y, Tong Y, Yu Z, Zyromski N, Schipani E (*I provided critical reagents and I critically reviewed the manuscript*) Carta F, Supuran CT, Korc M, Ivan M, Kelley MR and Fishel ML. Regulation of HIF1α under Hypoxia by APE1/Ref-1 Impacts CA9 Expression: Dual-Targeting in Patient-Derived 3D Pancreatic Cancer Models. Mol. Cancer Ther. 2016;15:2722-32
- 112. Agarwal S, Loder SJ, Cholok D, Peterson J, Li J, Breuler C, Cameron Brownley R, Hsin Sung H, Chung M, Kamiya N, Li S, Zhao B, Kaartinen V, Davis TA, Qureshi AT, Schipani E (*I provided critical reagents and I critically reviewed the manuscript*), Mishina Y, Levi B. Scleraxis-Lineage Cells Contribute to Ectopic Bone Formation in Muscle and Tendon. Stem Cells 2017; 35:705-10
- 113. Wang C, Xu CX, Alipppe Y, Qu C, Xiao J, **Schipani E** (*I provided critical reagents and I critically reviewed the manuscript*), Civitelli R, Abu-Amer Y and Mbalaviele G. Chronic inflammation triggered by NLRP3 inflammosome in myeloid cells promotes growth plate dysplasia by mesenchymal cells. **Sci.Rep. 2017**; 7:4880.
- 114. Clinkenbeard EL, Hanudel MR, Stayrook KR, Appaiah HN, Farrow EG, Cass TA, Summers LJ, Ip CS, Hum JM, Thomas JC, Ivan M, Richine BM, Chan RJ, Clemens TL, **Schipani E** (*I provided critical reagents, I contributed to the design of the study and I critically reviewed the*

manuscript), Sabbagh Y, Xu L, Srour EF, Alvarez MB, Kacena MA, Salusky IB, Nemeth E, White KE. Erythopoietin stimulates murine and human fibroblast growth factor-23, revealing novel roles for bone and bone marrow. **Haematologica 2017**; 102(11):e427-e43

- 115. Silagi ES, Schoepflin ZR, Seifert EL, Merceron C, Schipani E(*I provided critical reagents and I critically reviewed the manuscript*), Shapiro EM and Risbud MV. Bicarbonate recycling by HIF-1-dependent Carbonic Anhydrase isoforms 9 and 12 is critical in maintaining intracellular pH and viability of nucleus pulposus cells. J Bone Miner Res. 2018;33:338-355
- 116. Brunetti G, Faienza MF, Colaianni G, Gigante I, Oranger A, Pignataro P, Ingravallo G, Di Benedetto A, Bortolotti S, Di Comite M, Storlino G, Lippo L, Ward-Kavanagh L, Mori G, Reseland JE, Passeri G, Schipani E (*I critically reviewed the manuscript*), Tamada K, Ware CF, Colucci S, Grano M. Impairment of bone remodeling in LIGHT/TNFS14-Deficient mice. J Bone Miner Res. 2018;33:704-719
- 117. Faienza MF, Brunetti G, Sanesi L. Colaianni G, Celi M, Piacente L, D'Amato G, Schipani E (*I critically reviewed the manuscript*), Colucci S, Grano M. High irisin levels are associated with better glycemic control and bone health in children with Type 1 diabetes. Diabetes Res Clin Pract. 2018; 141:10-17
- 118. Liu Q, Wang J, Chen Y, Zhang Z, Saunders L, Schipani E (*I critically reviewed the manuscript*), Chen Q, Ma PX. Suppressing mesenchymal stem cell hypertrophy and endochondral ossification in 3D cartilage regeneration with nanofibrous poly(l-lactic acid) scaffold and matrilin 3. Acta Biomater. 2018;76:29-38
- 119. Colaianni G, Lippo L, Sanesi L, Brunetti G, Celi M, Cirulli N, Passeri G, Reseland J, Schipani E (*I critically reviewed the manuscript*), Faienza MF, Tarantino U, Colucci S, Grano M. Deletion of the Transcription Factor PGC-1α in Mice Negatively Regulates Bone Mass. Calcif Tissue Int. 2018; 103:638-652
- 120. Yao Q, Parvez Khan M, Merceron C, LaGory E, Tata Z, Mangiavini L, Hu J, Vemulapalli K, Chandel NS, Giaccia AJ, Schipani E. Suppressing mitochondrial respiration is critical for hypoxia tolerance in the fetal growth plate. Developmental Cell 201;49: 748-763 (Preview in the same issue of Dev Cell; featured in Science Signaling 2019;12). Oxygen (O₂) is both an indispensable metabolic substrate in various enzymatic reactions, including mitochondrial respiration, and a regulatory signal that controls the stability and activity of the transcription factor Hypoxia-Inducible Factor 1alpha (Hif1a), a key mediator of the cellular adaptation to low O₂ tension (hypoxia). Hypoxic cells require Hifla to survive. Additionally, Hifla is an inhibitor of mitochondrial respiration. Hence, we hypothesized that enhancing mitochondrial respiration is detrimental to the survival of hypoxic cells in vivo. We addressed this question in the fetal growth plate, which is an avascular and hypoxic structure formed by chondrocytes and a matrix enriched in collagens and proteoglycans. Our study demonstrates that mitochondrial respiration is dispensable for growth plate development during fetal life. More importantly, its impairment prevents both the extreme hypoxia as well as the pathological and massive cell death observed in fetal growth plates lacking Hifla. These findings show that enhancing mitochondrial respiration is detrimental

to the survival of growth plate chondrocytes by, at least in part, increasing intracellular hypoxia. We thus propose that partial suppression of mitochondrial respiration is crucial during normal embryonic development to protect the tissues that are physiologically hypoxic from lethal intracellular anoxia.

- 121. Merceron C, Ranganathan K, Castellini L, Wang E, Tata Z, Mangiavini L, Parvez Khan M, Levi B, Giaccia AJ, Schipani E. Hypoxia Inducible Factor-2alpha is a negative regulator of osteoblastogenesis and bone mass accrual. Bone Research 2019;7:7. Osteoblasts, which are the cells forming bone, operate in a hypoxic environment. Hypoxia Inducible Factor-2alpha (HIF2) is one of the mediators of the cellular response to hypoxia. However, its role in the control osteoblast biology is still poorly understood. In this study, we used mouse genetic and demonstrated that HIF2 is an inhibitor of osteoblastogenesis and bone mass accrual. Moreover, we provided evidence that HIF2 may impair osteoblast differentiation at least in part by upregulating the transcription factor Sox9. HIF2 can be selectively inhibited by small molecules that are currently in clinical trials in patients with renal carcinoma. Inhibiting HIF2 could thus represent a therapeutical approach for the treatment of the low bone mass observed in chronic diseases, osteoporosis or aging.
- 122. Noda H, Guo J, Khatri A, Dean T, Reyes M, Armanini M, Brooks DJ, Martins JS, Schipani E (*I provided critical animal models and reviewed the manuscript*), Bouxsein ML, Demay MB, Potts JT Jr, Jüppner H, Gardella TJ. An Inverse Agonist Ligand of the PTH Receptor Partially Rescues Skeletal Defects in a Mouse Model of Jansen's Metaphyseal Chondrodysplasia. J Bone Miner Res. 2020;35:540-549.
- 123. Yao Q, Parvez-Khan M, Schipani E. In vivo survival strategies for cellular adaptation to hypoxia: HIF1α-dependent suppression of mitochondrial oxygen consumption and decrease of intracellular hypoxia are critical for survival of hypoxic chondrocytes. Bone. 2020;140:115572.
- 124. Sartawi Z, Waeber C, **Schipani E** (*I contributed to the histolgical analysis and I critically reviewed the manuscript*), Ryan KB. Development of electrospun polymer scaffolds for the localized and controlled delivery of siponimod for the management of critical bone defects. **Int J Pharm. 2020** 30;590:119956.
- 125. Garcia-Galiano D, Cara AL, Tata Z, Allen SJ, Myers MG Jr, Schipani E (I contributed to the design of the study and the histological analysis, and I critically reviewed the manuscript), Elias CF. ERα Signaling in GHRH/Kiss1 Dual-Phenotype Neurons Plays Sex-Specific Roles in Growth and Puberty. J Neurosci. 2020;40:9455-9466.
- 126. Tata Z, Merceron C, **Schipani E**. Fetal Growth Plate Cartilage: Histological and Immunohistochemical Techniques. **Methods Mol Biol 2021**; 2254:53-84.
- 127. Silagi ES, **Schipani E**, Shapiro IM, Risbud MV. The role of HIF proteins in maintaining the metabolic health of the interevertebral disc. **Nat Rev Rheumatol 2021**;17:426-439.

- 128. Colucci SC, Buccoliero C, Sanesi L, Errede M, Colaianni G, Annese T, Khan MP, Zerlotin R, Dicarlo M, Schipani E (*I contributed to the design of the study and the histological analysis, and I critically reviewed the manuscript*), Kozloff KM, Grano M. Systemic Administration of Recombinant Irisin Accelerates Fracture Healing in Mice. Int J Mol Sci. 2021;22:10863.
- 129. Tolonen JP, Salo AM, Finnilä M, Aro E, Karjalainen E, Ronkainen VP, Drushinin K, Merceron C, Izzi V, Schipani E (*I contributed to the design of the study and the histological analysis, and I critically reviewed the manuscript*), Myllyharju J. Reduced Bone Mass in Collagen Prolyl 4-Hydroxylase *P4ha1*^{+/-}; *P4ha2*^{-/-} Compound Mutant Mice. JBMR Plus. 2022; 6(6):e10630.
- Mendoza-Reinoso V, Schnepp PM, Baek DY, Rubin JR, Schipani E (I contributed to the design of the study and I critically reviewed the manuscript), Keller ET, McCauley LK, Roca H. Bone marrow macrophages induce inflammation by efferocytosis of apoptotic prostate cancer cells via HIF-1alpha stabilization. Cells. 2022 ;11(23):3712.
- 131. Oranger A, Zerlotin R, Buccoliero C, Sanesi L, Storlino G, Schipani E (*I contributed to the design of the study and I critically reviewed the manuscript*), Kozloff KM, Mori G, Colaianni G, Colucci S, Grano M. Irisin modulates inflammatory, angiogenic, and osteogenic factors during fracture healing. International Journal of Molecular Sciences. 2023; 24(3): 1809. doi: 10.3390/ijms24031809.
- Lanzolla G, Khan MP, Sabini E, Giaccia A, Schipani E. Erytropoietin and skeletal crosstalks in physiology and disease. Current Opinion in Endocrine and Metabolic Research 2023; 29: 100436.
- 133. Sabini E, Arboit L, Khan MP, Lanzolla G, Schipani E. Oxidative phosphorylation in bone cells. Bone Rep. 2023;18:101688. doi: 10.1016/j.bonr.2023.101688.
- 134. Schipani E, Carmeliet G. Cell metabolism ad bone cells. Bone Rep. 2023;19:101719. doi: 10.1016/j.bonr.2023.101719
- 135. Warren A, Porter RM, Reyes-Castro O, Ali MM, Marques-Carvalho A, Kim HN, Gatrell LB, Schipani E (*I contributed to the design of the study and I critically reviewed the manuscript*), Nookaew I, O'Brien CA, Morello R, Almeida M. The NAD salvage pathway in mesenchymal cells is indisoensable for skeletal development in mice. Nat Commun. 2023;14(1):3616
- 136. Reyes M, Firat D, Hanna P, Khan M, Bruce M, Shvedova M, Kobayashi T, Schipani E (*I contributed to the design of the study and I critically reviewed the manuscript*), Gradella TJ, Jueppner H,. Substantially delayed maturation of growth plate chondrocytes in "humanized" PTH1R mice with the H223R mutation of Jansen's disease. JBMR Plus 2023; 7(10):e10802.dol 10.1002/jbm44.10802

- Sabini E, Schipani E. The hypoxia signature across skeletal progenitor cells. J Bone Miner Res. 2024;39(4):373-374
- 138. Lanzolla G, Merceron C, Khan MP, Sabini E, Giaccia A, **Schipani E**. Osteoblastic erythropoietin is not required for bone mass accrual. **JBMR Plus 2024**; 8(6). *Our study explores the role of erythropoietin (EPO), a hormone best known for its role in producing red blood cells, in bone health. While the kidneys produce most EPO, we now know that bone cells, specifically osteoblasts (which build bone), can also produce and release EPO when oxygen levels are low. However, the importance of this bone-derived EPO remains unclear.* To investigate, we created a mouse model where EPO production was specifically turned *off in bone-building cells and their precursors. In young adult mice, we found that the absence of osteoblast-derived EPO did not significantly impact bone growth or red blood cell production. Our findings suggest that EPO from osteoblasts may not play a major role in bone or blood health during early adulthood. However, we believe it could be more important under specific conditions, such as aging, bone healing, or disease. Future studies will help us better understand the potential functions of EPO produced by bone cells in these contexts.*
- 139. Khan MP, Sabini E, Beigel K, Lanzolla G, Laslow BM, Wang D, Merceron C, Giaccia A, Long F, Taylor DM, Schipani E. HIF1 safeguards cortical bone formation against impaired oxidative phosphorylation. JCI Insight 2024; 9(18):e182330. Our study explores how cells produce energy and how this affects the development of strong, healthy bones. Bones, especially the outer shell of long bones called cortical bone, need energy to grow and maintain their structure. Cells use two main ways to make energy: oxidative phosphorylation (OxPhos), which relies on oxygen and mitochondria (the cell's power plants), and glycolysis, which can work without oxygen. We focused on a protein called TFAM, which helps mitochondria function properly. In mice that were missing TFAM in certain bone cells, the outer shell of their long bones became thinner and weaker, leading to fractures. These mice also had problems with energy production in a layer of bone-forming cells called the periosteum, which is crucial for bone growth and repair. Interestingly, we found that activating another protein, HIF1, could help compensate for the energy production problems caused by TFAM loss. HIF1 pushes cells to use glycolysis instead of OxPhos. By boosting HIF1, some of the bone defects were improved, showing that cells can switch energy strategies to some extent. The significance of our study is twofold: it highlights how important mitochondria and OxPhos are for building strong bones, and it suggests that tweaking how cells make energy could be a future treatment for conditions like osteoporosis or other bone fragility diseases.
- 140. Lanzolla G, Sabini E, Beigel K, Khan MP, Sherry Liu X, Wang D, Laslow B, Taylor D, Bellido T, Giaccia A, Schipani E. Pharmacological inhibition of HIF2 protects against bone loss in an experimental model of estrogen deficiency. PNAS 2024;121. In this study, we looked at how a lack of estrogen, such as during menopause or certain medical conditions,

December 2024

can lead to weaker bones and a higher risk of fractures. Our focus was on the spongy part of bones called trabecular bone, which is particularly affected. Bone health depends on a balance between making new bone and breaking down old bone, and estrogen deficiency disrupts this balance. We investigated how HIF-2a (HIF2), which helps cells respond to low oxygen levels in the bone marrow, affects this process. We discovered that reducing or blocking HIF2 in bone-making cells during development leads to stronger bones with more trabecular bone mass. This happens because more bone-forming cells, called osteoblasts, are created. Excitingly, we tested a drug called PT2399 that blocks HIF2 and found it could prevent bone loss in mice with low estrogen levels, mimicking menopause. This drug increased the number of bone-forming cells by expanding the pool of early bone cell precursors.In short, our study highlights a new way to strengthen bones and prevent bone loss in conditions like menopause by targeting HIF2, offering potential for better treatments in the future.

Completed Publications in Scientific Journals: Not Peer-Reviewed

- 1. Schipani E, Bergwitz C, Kronenberg HM, Segre GV, Jüppner H. The human PTH/PTHrP receptor. In: DeBellis A, Schipani E, eds. Future Trends in Endocrinology, Frontiers in Endocrinology Series 1995; 14:15-20.
- 2. Schipani E. Natriuretic peptides, cGMP, and growth plate development. BoneKEy commentary, IBMS online 2001. (Invited)
- 3. Schipani E. A genetic dissection of IKKα functions. BoneKEy commentary, IBMS online 2002. (Invited)
- 4. Schipani E. A novel PTH/PTHrP receptor (PPRc) mutation: the ongoing tale of PPRc and the growth plate becomes more complex. BoneKEy commentary, IBMS online 2002. (Invited)
- 5. Schipani E. Puzzles of Cartilage Biology. In "Meeting Report from the 24th Annual Meeting of the American Society for Bone and Mineral Research." BoneKEy commentary, **IBMS** online 2002. (Invited)
- 6. Schipani E. Otoconin 22 and Calcitonin: A novel modality of regulating calcium storages in lower vertebrates? Endocrinology 2003, News and Views; 144:3285-3286. (Invited)
- Schipani E. Growth plate development: new pieces added to the puzzle. In: "Meeting Report from the 26th Annual Meeting of the American Society for Bone and Mineral Research." BoneKEy commentary, IBMS online 2004. (Invited)
- 8. Schipani E. Mutations of preproparathyroid hormone gene in primary hyperparathyroidism. Clinical Cases in Mineral and Bone Metabolism 2004; 1:107-108. (Invited)
- 9. Schipani E. On the road to the "big" chondrocytes. In: "Meeting Report from the 27th Annual Meeting of the American Society for Bone and Mineral Research." BoneKEy commentary, **IBMS online 2005**. (Invited)

- 10. Schipani E. Chondrocytes: old friends and new acquaintances. In: "Meeting Report from the 28th Annual Meeting of the American Society for Bone and Mineral Research." BoneKEy commentary, **IBMS online 2006**. (Invited)
- 11. Schipani E. Chondrocytes: a few pearls in an ocean of bones. In "Meeting Report from the 29th Annual Meeting of the American Society for Bone and Mineral Research." BoneKEy commentary, **IBMS online 2007.** (Invited)
- 12. Schipani E, Clemens TL: Hypoxia and the Hypoxia-Inducible Factors in the Skeleton. BoneKey 2008; 5:275-284. (Invited)
- 13. Merceron C. and **Schipani E**. Chondrocytes: a few grains of softness in a hard bone world. **BoneKey 2013** (Meeting Report of the Annual Meeting of the American Society for Bone and Mineral Research 2012. (Invited)
- Merceron C. and Schipani E. Intervertebral Disc: a rising star in the skeleton galaxy. BoneKey 2013 Meeting Report of the Annual Meeting of the American Society for Bone and Mineral Research 2012. (Invited)
- 15. Schipani E and Kobayashi T. Editorial for "Special Issue on the Growth Plate". Bone 2020; 144:115796

Books

- 1. Bilezikian JP, Yoshimoto K, Pollak MR, Schipani E, Jüppner H, Brown EM, Arnold A, Thakker RV. Genetic Disorders of Parathyroid Hormone Action. In "Genetics in Endocrinology," Lippincott, 2002.
- 2. Carter PH, Schipani E. The PTH/PTHrP receptor in human diseases. In "Hormone Resistance and Hypersensitivity States." Eds GP Chrousos, JM Olefsky, E Samols, Lippincott Williams and Wilkins, 2002, p269-288. (Invited)
- Jüppner H, Schipani E, Silve C. Jansen's metaphyseal chondrodysplasia and Blomstrand's lethal chondrodysplasia: two genetic disorders caused by PTH/PTHrP receptor mutations. In "Principles of Bone Biology", Eds JP Bilezikian, LG Raisz, GA Rodan. 2nd edition, Academic Press, 2002, p1117-1135.
- 4. Schipani E. Jansen's metaphyseal chondrodysplasia. In "NORD Guide to Rare Disorders", Ed Ed Gruson, Lippincott, 2003, pp 206-207.
- Provot S, Schipani E, Wu J, Kronenberg H. Development of the Skeleton. In "Osteoporosis", Eds R Marcus, D Feldman, DA Nelson, CJ Rosen. 3rd edition, Academic Press, 2007, p241-269. (Invited)
- 6. Schipani E, Khatri R. The role of hypoxia-induced factors. In "Bone and Development" Eds Bronner F, Farach-Carson M, Roach HI; Springer "Topics in Bone Biology" Series, 2010; 107-123. (Invited)
- 7. Mangiavini L, O'Riley R and Schipani E. Use of Knockout and Transgenic Mouse Models in Disc Research. In "Intervertebral Disc" Eds M.Risbud and I.Shapiro, 2013, p 341-352. (Invited)

- 8. Mangiavini L and Schipani E. TUNEL assay for detection of cell death. Methods Mol Biol 2014; 1130:245-8. (Invited)
- 9. Mangiavini L, Merceron C and Schipani E. Analysis of mouse growth plate development. Current Protocols in Mouse Biology 2016; 6: 67-130. (Invited)
- 10. Tata Z, Merceron C, Schipani E. Fetal Growth Plate Cartilage: Histological and Immunohistochemical Techniques. Methods Mol Biol. 2021;2245:53-84. (Invited)

Other media

1. McMahon G, Schipani E, Strewler G. Tutorial case and tutor guide, "An appetite for success." Disorders of the Musculoskeletal, Endocrine and Reproductive Systems. Human Systems. Harvard Medical School, 2004.