

This article was downloaded by: [University of Pennsylvania]

On: 15 February 2013, At: 06:26

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



AIDS Care: Psychological and Socio-medical Aspects of AIDS/HIV

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/caic20>

Effects of behavioral stress reduction Transcendental Meditation intervention in persons with HIV

Sumedha Chhatre ^a, David S. Metzger ^a, Ian Frank ^b, Jean Boyer ^c, Edward Thompson ^c, Sanford Nidich ^d, Luis J. Montaner ^e & Ravishankar Jayadevappa ^f

^a Department of Psychiatry, HIV/AIDS Prevention Research Division, University of Pennsylvania, Philadelphia, PA, USA

^b Department of Medicine, Infectious Disease Division, University of Pennsylvania, Philadelphia, PA, USA

^c Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA, USA

^d Department of Physiology and Health, Maharishi University of Management, Fairfield, IA, USA

^e Wistar Institute, Philadelphia, PA, USA

^f Department of Medicine, Geriatrics Division, University of Pennsylvania, PA, USA

Version of record first published: 11 Feb 2013.

To cite this article: Sumedha Chhatre, David S. Metzger, Ian Frank, Jean Boyer, Edward Thompson, Sanford Nidich, Luis J. Montaner & Ravishankar Jayadevappa (2013): Effects of behavioral stress reduction Transcendental Meditation intervention in persons with HIV, *AIDS Care: Psychological and Socio-medical Aspects of AIDS/HIV*, DOI:10.1080/09540121.2013.764396

To link to this article: <http://dx.doi.org/10.1080/09540121.2013.764396>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Effects of behavioral stress reduction Transcendental Meditation intervention in persons with HIV

Sumedha Chhatre^{a*}, David S. Metzger^a, Ian Frank^b, Jean Boyer^c, Edward Thompson^c, Sanford Nidich^d, Luis J. Montaner^e and Ravishankar Jayadevappa^f

^aDepartment of Psychiatry, HIV/AIDS Prevention Research Division, University of Pennsylvania, Philadelphia, PA, USA; ^bDepartment of Medicine, Infectious Disease Division, University of Pennsylvania, Philadelphia, PA, USA; ^cDepartment of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA, USA; ^dDepartment of Physiology and Health, Maharishi University of Management, Fairfield, IA, USA; ^eWistar Institute, Philadelphia, PA, USA; ^fDepartment of Medicine, Geriatrics Division, University of Pennsylvania, PA, USA

(Received 30 September 2012; final version received 1 January 2013)

Stress is implicated in the pathogenesis and progression of HIV. The Transcendental Meditation (TM) is a behavioral stress reduction program that incorporates mind–body approach, and has demonstrated effectiveness in improving outcomes via stress reduction. We evaluated the feasibility of implementing TM and its effects on outcomes in persons with HIV. In this community-based single blinded Phase-I, randomized controlled trial, outcomes (psychological and physiological stress, immune activation, generic and HIV-specific health-related quality of life, depression and quality of well-being) were assessed at baseline and at six months, and were compared using parametric and nonparametric tests. Twenty-two persons with HIV were equally randomized to TM intervention or healthy eating (HE) education control group. Retention was 100% in TM group and 91% in HE control group. The TM group exhibited significant improvement in vitality. Significant between group differences were observed for generic and HIV-specific health-related quality of life. Small sample size may possibly limit the ability to observe significant differences in some outcomes. TM stress reduction intervention in community dwelling adults with HIV is viable and can enhance health-related quality of life. Further research with large sample and longer follow-up is needed to validate our results.

Keywords: HIV infection; stress reduction; Transcendental Meditation; health-related quality of life; immune activation

Introduction

The number of HIV/AIDS survivors in the USA has increased from 16,8754 to 48,7968 between 1993 and 2009 (CDC, 2012). Improved survival implies that persons with HIV are vulnerable to HIV-specific and non-HIV-specific conditions such as immune activation and impaired well-being (Desai & Landay, 2010; Kovacs et al., 2010).

Living with HIV involves facing psychological and physical stressors. Relationship between stress, HIV, and health is complex (Bonneau, Padgett, & Sheridan, 2007; Cohen, Janicki-Deverts, & Miller, 2007; Fumaz et al., 2009). Studies have explored the association between stress management and psychosocial outcomes in HIV (Carrico et al., 2006; Leserman, 2008; Robinson, Mathews, & Witek-Janusek, 2003). Cortisol is used as a measure of physiological stress (Cruess, Antoni, Kumar, & Schneiderman, 2000; Cruess et al., 2000; Goodkin et al., 1996; Ironson et al., 1996; Leserman et al., 2002). Some studies indicate improved immunological function for stress management groups (Antoni et al., 1991; Creswell, Myers,

Cole, & Irwin, 2009; Robinson et al., 2003), while others report improvements for intervention and control groups, or no effect (Coates, McKusick, Kuno, & Stites, 1989; LaPerriere et al., 1990).

Psychoneuroimmunology (PNI) theory provides a framework for analyzing stress–disease relationships (Hand, Phillips, Dudgeon, & Skelton, 2005). Transcendental Meditation (TM) is a behavioral stress-reduction program that incorporates mind–body approach. Studies suggest that TM may improve outcomes across various illnesses (Castillo-Richmond et al., 2000; Jayadevappa et al., 2007; Rainforth et al., 2007; Schneider et al., 2001). Objective of our Phase-I study was to evaluate the feasibility of TM intervention in persons with HIV and analyze the potential effects of stress reduction on psychosocial, hormonal, and immunological outcomes.

Methods

This two arm single blinded randomized control trial (RCT) was approved by the Institutional Review

*Corresponding author. Email: rasu@mail.med.upenn.edu

Board. All participants provided written consent. Persons with HIV satisfying following criteria were eligible: age ≥ 18 years; stable on antiretroviral treatment; CD4+ T-cell count > 300 and plasma viral load of < 200 copies/ml in the past six months; no Hepatitis C, non-diabetic; not on steroids; minimal test score of ≥ 25 ; and not participating in other studies. Intervention group underwent the TM program of six-month duration. Control group underwent a healthy eating (HE) education program of comparable length and structure. We used a community-based recruitment initiative. Between April 8 and May 10, 2011, 50 persons contacted the study team, 34 underwent eligibility assessment, and 22 were randomized after baseline measurements (Figure 1).

The TM intervention was conducted by a certified instructor and comprised of two phases. Intensive phase consisted of five consecutive sessions of two-hour duration each day. Follow-up phase involved bi-weekly sessions for three months, followed by monthly sessions for additional three months. Participants were asked to practice TM twice a day for 20 minutes each over the study period.

The HE education control program was designed to mirror the effects of personal interaction and attention that are part of TM program. This course was conducted by a nurse and addressed more in-depth nutritional information than generally is provided in HIV care. Similar to TM intervention, intensive phase of HE program consisted of five consecutive sessions of two-hour duration each day. Follow-up phase involved bi-weekly sessions for three months, followed by monthly sessions for three additional months, and were conducted by the same nurse. Participants were asked to engage in an activity, e.g., reading, twice a day for 20 minutes during the study.

Outcomes

Outcomes were assessed at baseline and at six month. We conceptualized feasibility as response to recruitment, and acceptability as study compliance.

Endocrine response was measured by serum norepinephrine and cortisol levels. Blood draws were done between 9 and 10 am and 25 μ l of sera was used to quantify norepinephrine levels (BA E-5200 Rocky Mountain Diagnostics, 2012). Cortisol




Figure 1. Recruitment flow diagram.

Table 1. Demographics of the study population.

	Transcendental Meditation (<i>n</i> = 11)	Healthy eating (HE) (<i>n</i> = 11)
Age in years (mean \pm std)	49.7 \pm 7.1	50.0 \pm 4.4
Gender male (%)	81.8	81.8
Race (%)		
African-American	63.6	81.8
White	9.1	18.2
Hispanic	27.3	0.00
Employment (%)		
Working (Full time/part time)	18.18	27.27
Other (retired, disabled)	81.82	72.73
Ever smoked (%)	54.6	50.00
Length of infection in years (mean \pm std)	16.5 \pm 6.8	11.2 \pm 6.2

levels were quantified using assay from Parameter Cortisol Assay (R & D Systems, 2011). Immune activation was measured by expression of CD38 and HLA-DR on CD4 and CD8 T cells and was analyzed using FlowJo (Treestar, 2011).

Table 2. Comparison of outcomes.

Variable (mean \pm standard deviation)	Transcendental Meditation <i>n</i> = 11		HE education <i>n</i> = 9		Change	
	Baseline	Six month	Baseline	Six month	TM change	HE change
Hormonal outcomes						
Cortisol (ng/ml)	2.36 \pm 1.30	2.21 \pm 1.65	2.59 \pm 1.17	3.01 \pm 0.87	0.14 \pm 1.17	-0.41 \pm 1.89
Norepinephrine (ng/ml)	1.05 \pm 0.31	0.91 \pm 0.58	0.93 \pm 0.20	0.94 \pm 0.48	0.13 \pm 0.64	-0.01 \pm 0.44
Psychological outcomes						
PSS	19.6 \pm 3.4	18.4 \pm 5.0	18.6 \pm 4.9	18.9 \pm 3.9	1.27 \pm 4.3	-0.33 \pm 0.61
CESD	18.4 \pm 9.2	13.5 \pm 11.5	19.7 \pm 7.7	20.1 \pm 15.1	4.8 \pm 10.8	-0.44 \pm 11.7
Health outcomes						
SF-36						
Physical function	60.9 \pm 15.14	55.5 \pm 24.84	48.3 \pm 26.69	41.1 \pm 23.42	5.45 \pm 26.68	7.2 \pm 17.52
Role physical	75.0 \pm 40.31	70.5 \pm 41.56	50.0 \pm 41.46	33.3 \pm 33.07	4.55 \pm 57.89	16.7 \pm 48.41
Role emotional	72.7 \pm 41.68	75.8 \pm 42.40	62.9 \pm 35.14	40.7 \pm 49.38	-3.03 \pm 64.04	22.2 \pm 47.14
Vitality	46.0 \pm 19.42	69.9 \pm 19.13*	48.6 \pm 25.15	52.8 \pm 25.98	-23.86 \pm 26.78	-4.17 \pm 24.41
Mental health	60.9 \pm 15.14	74.1 \pm 19.98	63.3 \pm 18.54	71.1 \pm 22.61	-13.18 \pm 27.95	-7.78 \pm 27.51
Social function	60.2 \pm 23.59	73.9 \pm 24.66	63.9 \pm 27.56	59.7 \pm 24.03	-13.04 \pm 33.75	4.17 \pm 30.62
Bodily pain	75.7 \pm 24.88	72.0 \pm 35.05	57.2 \pm 27.85	67.2 \pm 28.32	3.63 \pm 34.34	-10.0 \pm 21.10
General health	51.4 \pm 21.34	66.8 \pm 18.48	70.0 \pm 20.92	63.9 \pm 16.16	-15.45 \pm 20.67	6.11 \pm 16.16*
FAHI						
Physical well-being	29.2 \pm 7.2	36.2 \pm 4.2*	28.1 \pm 0.23	29.3 \pm 7.9	-7.0 \pm 6.9	-1.2 \pm 4.6
Emotional well-being	28.1 \pm 8.9	34.3 \pm 6.0	25.7 \pm 6.71	25.7 \pm 10.6	-6.8 \pm 6.6	0 \pm 10.0
Functional and global well-being	34.6 \pm 11.2	38.1 \pm 10.1	34.6 \pm 10.26	33.0 \pm 12.1	-3.4 \pm 4.8	1.6 \pm 5.4
Social well-being	22.4 \pm 7.4	23.0 \pm 9.3	18.7 \pm 8.51	18.6 \pm 8.1	-0.63 \pm 3.5	0.11 \pm 6.7
Cognitive functioning	8.7 \pm 2.9	9.3 \pm 2.9	8.6 \pm 2.35	7.9 \pm 2.8	-0.54 \pm 3.2	0.7 \pm 3.3
Total score	123.0 \pm 28.2	140.8 \pm 26.5	115.6 \pm 25.27	114.4 \pm 29.1	-17.8 \pm 17.9	1.1 \pm 19.4*
Quality of well-being	0.69 \pm 0.19	0.72 \pm 0.22	0.60 \pm 0.22	0.50 \pm 0.19	-0.03 \pm 0.27	0.09 \pm 0.23

Note: *Significant at *p* < 0.05.

Perceived Stress Scale (PSS) was used to measure psychological stress (Cohen & Williamson, 1988). Generic health-related quality of life (HRQoL) was measured using Medical Outcomes Short Form, SF-36 (Ware & Sherbourne, 1992) and HIV-specific HRQoL was measured using Functional Assessment of HIV Infection (FAHI) instrument (Peterman, Cella, Mo, & McCain, 1997). Center for Epidemiological Studies Depression (CESD) scale was used to measure depression (Zich, Attkisson, & Greenfield, 1990). Quality of well-being was measured using Quality of Well-being (QWB-SA) survey (Kaplan & Bush, 1982).

Analysis

Data were analyzed using intent-to-treat approach. We compared distributions of socio-demographic variables between the groups to determine the effectiveness of randomization. Due to small sample size and skewed distribution for some outcomes, we log transformed the outcome scores and performed paired *t*-tests for within group changes. For between group effects, changes in scores from baseline to six month were tested using independent *t*-tests.




Figure 2. Change in immune activation markers.

Results

Retention was 100% for TM group and 91% for control group, indicating good feasibility and acceptability of the intervention. Socio-demographic characteristics were comparable between groups (Table 1). Results of within-and-between group analyses are presented in Table 2. At follow-up, perceived stress in TM group was lower than baseline and remained mostly unchanged in the controls. Mean baseline cortisol was comparable between groups. Cortisol was somewhat lower in TM group at follow-up, and higher in the controls. Norepinephrine declined slightly in TM group, and remained unchanged in the control group. While there was no significant difference in level of T cell activation between groups, differences in TM group were closer to zero compared to the controls (Figure 2). Additionally, a two-part test comparing changes in T cell activation between groups found lower change in HD Effectors CD8+ HLA-DR+CD38+freq of CD8+T cells for TM group ($p=0.0633$), indicating possibility for immune activation stabilization.

For TM group, SF-36 domains of role emotional, vitality ($p=0.0133$), mental health, social function, and general health showed improvement. General health improved in TM group, and declined in the controls ($p=0.0299$). Among domains of FAHI, improvement in physical well-being was larger for TM group compared to controls ($p=0.0127$). Between-group changes were significant for FAHI total score ($p=0.0393$). Mean CESD score decreased in TM group and remained unchanged in the controls. The QWB improved in TM group, and declined in the controls.

Discussion

In this Phase-I, community-based RCT we observed the feasibility and acceptability of TM stress-reduction program in persons with HIV. The TM group exhibited significant improvement in generic and HIV-specific HRQoL. We noticed a pattern of improvement in physiological and psychological parameters in TM group. Some differences were

statistically nonsignificant, possibly due to small sample size and short follow-up.

Research using PNI framework reports benefits of stress reduction in HIV patients (Antoni et al., 1991; Coates et al., 1989; Ironson et al., 1996; Lutgendorf et al., 1997; McCain & Zelle, 1996). Stress reduction was associated with reduction in depression (Antoni et al., 1991; Lutgendorf et al., 1997), anxiety (Ironson et al., 1996; Lutgendorf et al., 1997), improvement in CD4 count (Antoni et al., 1991), Natural Killer (NK) cell number (Antoni et al., 1991), safer sexual practice (Coates et al., 1989), and emotional well-being (McCain & Zelle, 1996).

Per PNI theory, we conceptualized the pathway where psychosocial/psychological factors are immediately affected by stress. This may be one of the reasons why changes in generic and HIV-specific HRQoL and perceived stress were observed in TM group. Some studies have reported significant effects of stress reduction on cortisol (D. G. Cruess et al., 2000; S. Cruess et al., 2000; Ironson et al., 1996). However, our results are similar to studies that reported no significant change in cortisol (Leserman et al., 2002; McCain et al., 2003; Robinson et al., 2003). Reasons for differential findings may be sample size, length of follow-up, type of sample (serum, salivary, or urine), time of measurement, and length of HIV-infection.

This is the first study to explore application of TM in community-dwelling adults with HIV and adds to the existing evidence regarding association between stress and outcomes. Unique features of our study are RCT design and assessment across all domains of PNI framework. We note following study limitations. Small sample size may possibly limit the ability to observe significant differences in some outcomes. Also, data about mental health treatment or other stress reducing activities of participants were not collected. Finally, those who participated may be more committed to behavior change, leading to self-selection bias.

Increasing numbers of HIV survivors are facing psychological and physical stressors with debilitating effects on health outcomes. Thus, addressing stress reduction in persons with HIV is critical. Our research shows that behavioral stress reduction TM intervention was acceptable in persons with HIV and improved generic and HIV-specific HRQoL. Future research should validate our findings, and help establish TM in particular, and stress reduction in general, as a complementary technique in the battle against HIV.

Acknowledgements

This pilot was funded by the University of Pennsylvania's Center for AIDS Research (CFAR) as part of the Developmental Award Program. We recognize the crucial support and commitment of our study participants. We thank Debora Dunbar, Sara Whittington, Dana Brown, Annet Davis-Vogel, Sheryl Foster, Kevin T. Jones, Chelsea Voytek, Danielle Fiore, Mack Taylor, Pete Smith, Tiffany B. Dominique, Ann Colon, LaRue Rainone and Robert Frederick of HIV Prevention Research Division, University of Pennsylvania; Glenn Davis, Transcendental Meditation instructor; and Julia Bohinski, Health Eating education instructor.

References

- Antoni, M. H., Baggett, L., Ironson, G., La Perriere, A., August, S., Klimas, N., ... Fletcher, M. A. (1991). Cognitive-behavioral stress management intervention buffers distress response and immunological changes following notification of HIV-1 seropositivity. *Journal of Consulting and Clinical Psychology, 59*, 906–915. doi:10.1037/0022-006X.59.6.906
- BA E-5200 Rocky Mountain Diagnostics. (2012). Colorado Springs, CO, USA.
- Bonneau, R. H., Padgett, D. A., & Sheridan, J. F. (2007). Twenty years of psychoneuroimmunology and viral infections in Brain, Behavior, and Immunity. *Brain Behavior and Immunity, 21*, 273–280. doi:10.1016/j.bbi.2006.10.004
- Carrico, A. W., Antoni, M. H., Duran, R. E., Ironson, G., Penedo, F., Fletcher, M. A., ... Schneiderman, N. (2006). Reductions in depressed mood and denial coping during cognitive behavioral stress management with HIV-positive gay men treated with HAART. *Annals of Behavioral Medicine, 31*, 155–164. doi:10.1207/s15324796abm3102_7
- Castillo-Richmond, A., Schneider, R. H., Alexander, C. N., Cook, R., Myers, H., Nidich, S., ... Salerno, J. (2000). Effects of the Transcendental Meditation Program on carotid atherosclerosis in hypertensive African Americans. *Stroke, 31*, 568–573. doi:10.1161/01.STR.31.3.568
- CDC. (2012). Retrieved from <http://www.cdc.gov/hiv/resources/factsheets/us.htm>
- Coates, T. J., McKusick, L., Kuno, R., & Stites, D. P. (1989). Stress reduction training changes number of sexual partners but not immune function in men with HIV. *American Journal of Public Health, 79*, 885–887. doi:10.2105/AJPH.79.7.885
- Cohen, S., & Williamson, A. (1988). Perceived stress in a probability sample of the United States. In S. Spacapan & S. Oskamp (Eds.), *Social Psychology of Health* (pp. 31–67). Newbury Park, CA: Sage.
- Cohen, S., Janicki-Deverts, D., & Miller, G. E. (2007). Psychological stress and disease. *JAMA: The Journal of the American Medical Association, 298*, 1685–1687. doi:10.1001/jama.298.14.1685

- Creswell, D. J., Myers, H. F., Cole, S. W., & Irwin, M. R. (2009). Mindfulness meditation training effects on CD4+ T lymphocytes in HIV-1 infected adults: A small randomized controlled trial. *Brain, Behavior, and Immunity*, *23*, 184–188. doi:10.1016/j.bbi.2008.07.004
- Cruess, D. G., Antoni, M. H., Kumar, M., & Schneiderman, N. (2000). Reductions in salivary cortisol are associated with mood improvement during relaxation training among HIV-seropositive men. *Journal of Behavioral Medicine*, *23*, 107–122. doi:10.1023/A:1005419917023
- Cruess, S., Antoni, M., Cruess, D., Fletcher, M. A., Ironson, G., Kumar, M., ... Schneiderman, N. (2000). Reductions in herpes simplex virus type 2 antibody titers after cognitive behavioral stress management and relationships with neuroendocrine function, relaxation skills, and social support in HIV-positive men. *Psychosomatic Medicine*, *62*, 828–837.
- Desai, S., & Landay, A. (2010). Early immune senescence in HIV disease. *Current HIV/AIDS Reports*, *7*, 4–10. doi:10.1007/s11904-009-0038-4
- Fumaz, C., Gonzalez-Garcia, M., Borrás, X., Ferrer, M. J., Muñoz-Moreno, J. A., Peña, R., ... Clotet, B. (2009). Increased peripheral proinflammatory cytokines in HIV-1-infected patients with prolonged viral suppression suffering from high psychological stress. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, *52*, 427–428. doi:10.1097/QAI.0b013e3181bb26a5
- Goodkin, K., Feaster, D. J., Tuttle, R., Blaney, N. T., Kumar, M., Baum, M. K., ... Fletcher, M. A. (1996). Bereavement is associated with time-dependent decrements in cellular immune function in asymptomatic human immunodeficiency virus type 1-seropositive homosexual men. *Clinical and Diagnostic Laboratory Immunology*, *3*, 109–118.
- Hand, G. A., Phillips, K. D., Dudgeon, W. D., & Skelton, W. D. (2005). Stress reduction as a means to enhance oral immunity in HIV infected individuals. *Journal of the Association of Nurses in AIDS Care*, *16*, 58–63. doi:10.1016/j.jana.2005.07.006
- Ironson, G., Field, T., Scafidi, F., Hashimoto, M., Kumar, M., Kumar, A., ... Fletcher, M. A. (1996). Massage therapy is associated with enhancement of the immune system's cytotoxic capacity. *International Journal of Neuroscience*, *84*, 205–217. doi:10.3109/00207459608987266
- Jayadevappa, R., Johnson, J. C., Bloom, B. S., Nidich, S., Desai, S., Chhatre, S., ... Schneider, R. (2007). Effectiveness of stress reduction program – Randomized pilot study. *Ethnicity and Disease*, *17*, 72–77.
- Kaplan, R. M., & Bush, W. (1982). Health-related quality of life measurement for evaluation research and policy analysis. *Health Psychology*, *1*, 61–80. doi:10.1037/0278-6133.1.1.61
- Kovacs, A., Karim, R., Mack, W. J., Xu, J., Chen, A., Operskalski, E., ... Al-Harhi, L. (2010). Activation of CD8 T cells predicts progression of HIV infection in women co infected with hepatitis C virus. *Journal of Infectious Diseases*, *201*, 823–834. doi:10.1086/650997
- LaPerriere, A. R., Antoni, M. H., Schneiderman, N., Ironson, G., Klimas, N., Caralis, P., & Fletcher, M. A. (1990). Exercise intervention attenuates emotional distress and natural killer cell decrements following notification of positive serological status for HIV-1. *Biofeedback and Self-Regulation*, *15*, 299–242. doi:10.1007/BF01011107
- Leserman, J. (2008). Role of depression, stress and trauma in HIV disease progression. *Psychosomatic Medicine*, *70*, 539–545. doi:10.1097/PSY.0b013e3181777a5f
- Leserman, J., Petitto, J. M., Gu, H., Gaynes, B. N., Barroso, J., Golden, R. N., ... Evans, D. L. (2002). Progression to AIDS, a clinical AIDS condition, and mortality: Psychosocial and physiological predictors. *Psychological Medicine*, *32*, 1059–1073. doi:10.1017/S0033291702005949
- Lutgendorf, S. K., Antoni, M. H., Ironson, G., Klimas, N., Kumar, M., Starr, K., ... Schneiderman, N. (1997). Cognitive-behavioral stress management decreases dysphoric mood and herpes simplex virus-type 2 antibody titres in symptomatic gay men. *Journal of Consulting and Clinical Psychology*, *65*, 31–43. doi:10.1037/0022-006X.65.1.31
- McCain, N. L., & Zelle, J. M. (1996). Psychoneuroimmunological studies in HIV disease. *Annual Review of Nursing Research*, *14*, 23–55.
- McCain, N. L., Munjas, B. A., Munro, C. L., Elswick, R.K., Robins, J. L., Ferreira-Gonzalez, A., ... Cochran, K. L. (2003). Effects of stress management on PNI-based outcomes in persons with HIV disease. *Research in Nursing and Health*, *26*(2), 102–117. doi:10.1002/nur.10074
- Peterman, A. H., Cella, D., Mo, F., & McCain, N. (1997). Psychometric validation of the revised Functional Assessment of Human Immunodeficiency Virus Infection (FAHI) quality of life instrument. *Quality of Life Research*, *6*, 572–584. doi:10.1023/A:1018416317546
- R & D Systems. (2011). Minneapolis, MN, USA.
- Rainforth, M. V., Schneider, R. H., Nidich, S. I., Gaylord-King, C., Salerno, J. W., & Anderson, J. W. (2007). Stress reduction programs in patients with elevated blood pressure: A systematic review and meta-analysis. *Current Hypertension Reports*, *9*, 520–528. doi:10.1007/s11906-007-0094-3
- Robinson, F., Mathews, H. L., & Witek-Janusek, L. (2003). Psycho-endocrine-immune responses to mindfulness-based stress reduction in individuals infected with the human immunodeficiency virus: A quasi-experimental study. *The Journal of Alternative and Complementary Medicine*, *9*, 683–694. doi:10.1089/107555303322524535
- Schneider, R. H., Castillo-Richmond, A., Alexander, C. N., Myers, H., Kaushik, V., Aranguri, C., ... Nidich, S. (2001). Behavioral treatment of hypertensive heart disease in African Americans: Rationale and design of a RCT. *Behavioral Medicine*, *27*, 83–95. doi:10.1080/08964280109595775
- Treestar Inc., (2011). Ashland, OR, USA.

- Ware, J. E., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36): Conceptual framework and item selection. *Medical Care*, *30*(6), 473–483. doi:10.1097/00005650-199206000-00002
- Zich, J. M., Attkisson, C. C., & Greenfield, T. K. (1990). Screening for depression in primary care clinics: The

CES-D and the BDI. *International Journal of Psychiatry in Medicine*, *20*, 259–277. doi:10.2190/LYKR-7VHP-YJEM-MKM2